

Ion Storage Ring Measurements of Low Temperature Dielectronic Recombination Rate Coefficients for Modeling X-Ray Photoionized Cosmic Plasmas

D. W. Savin

*Columbia Astrophysics Laboratory and Department of Physics,
Columbia University, New York, NY*

G. Gwinner, D. Schwalm, and A. Wolf

*Max-Planck-Institut für Kernphysik, Heidelberg, Germany
and Physikalisches Institut der Universität Heidelberg, Heidelberg, Germany*

A. Müller and S. Schippers

Institut für Kernphysik, Strahlencentrum der Justus-Liebig-Universität, Giessen, Germany

1. Reliable DR Data Needed

Low temperature dielectronic recombination (DR) is the dominant recombination mechanism for most ions in X-ray photoionized cosmic plasmas. Reliably modeling and interpreting spectra from these plasmas requires accurate low temperature DR rate coefficients. Of particular importance are the DR rate coefficients for the iron L -shell ions (Fe XVII-Fe XXIV). These ions are predicted to play an important role in determining the thermal structure and line emission of X-ray photoionized plasmas, which form in the media surrounding accretion powered sources such as X-ray binaries (XRBs), active galactic nuclei (AGN), and cataclysmic variables (Savin *et al.*, 2000).

The need for reliable DR data of iron L -shell ions has become particularly urgent after the launches of *Chandra* and *XMM-Newton*. These satellites are now providing high-resolution X-ray spectra from a wide range of X-ray photoionized sources. Interpreting the spectra from these sources requires reliable DR rate coefficients. However, at the temperatures relevant for X-ray photoionized plasmas, existing theoretical DR rate coefficients can differ from one another by factors of two to orders of magnitudes.

2. New Program at Max-Planck-Institute

To address the need for accurate low temperature DR rate coefficients of the iron L -shell ions, we have initiated a program of measurements for DR via $2 \rightarrow 2$ core excitations using the heavy-ion Test Storage Ring located at the Max-Planck-Institute for Nuclear Physics in Heidelberg, Germany. To date measurements have been carried out for Fe XVIII (Savin *et al.*, 1997, 1999), Fe XIX (Savin *et al.*, 1999), Fe XX (Savin *et al.* 2002), Fe XXI, Fe XXII, and Fe XXIV. A detailed discussion of our results can be found in the references cited.

Acknowledgments

This work was supported in part by NASA SARA Program grant NAG5-5261, the German Federal Minister for Education and Research (BMBF), and the German Research Council (DFG).

REFERENCES

- Savin, D. W., *et al.*, 1997, *ApJ*, **489**, L115.
Savin, D. W., *et al.*, 1999, *ApJS*, **123**, 687.
Savin, D. W., *et al.*, 2000, in *Proceedings of the 12th APS Topical Conference on Atomic Processes in Plasmas*, Reno Nevada, ed. R. C. Mancini (New York: American Institute of Physics), p. 267.
Savin, D. W., *et al.*, 2002, *ApJS*, **138**, 337.