Complex effects of inhibiting hepatic apolipoprotein B100 synthesis in humans

Gisette Reyes-Soffer,^{1*} Byoung Moon,¹ Antonio Hernandez-Ono,¹ Marija Dionizovik-Dimanovski,¹ Jhonsua Jimenez,¹ Joseph Obunike,² Tiffany Thomas,¹ Colleen Ngai,¹ Nelson Fontanez,¹ Daniel S. Donovan,¹ Wahida Karmally,¹ Stephen Holleran,¹ Rajasekhar Ramakrishnan,¹ Robert S. Mittleman,^{3†} Henry N. Ginsberg^{1*}

Mipomersen is a 20mer antisense oligonucleotide (ASO) that inhibits apolipoprotein B (apoB) synthesis; its low-density lipoprotein (LDL)-lowering effects should therefore result from reduced secretion of very-low-density lipoprotein (VLDL). We enrolled 17 healthy volunteers who received placebo injections weekly for 3 weeks followed by mipomersen weekly for 7 to 9 weeks. Stable isotopes were used after each treatment to determine fractional catabolic rates and production rates of apoB in VLDL, IDL (intermediate-density lipoprotein), and LDL, and of triglycerides in VLDL. Mipomersen significantly reduced apoB in VLDL, IDL, and LDL, which was associated with increases in fractional catabolic rates of VLDL and LDL apoB and reductions in production rates of IDL and LDL apoB. Unexpectedly, the production rates of VLDL apoB and VLDL triglycerides were unaffected. Small interfering RNA-mediated knockdown of apoB expression in human liver cells demonstrated preservation of apoB secretion across a range of apoB synthesis. Titrated ASO knockdown of *apoB* mRNA in chow-fed mice preserved both apoB and triglyceride secretion. In contrast, titrated ASO knockdown of *apoB* mRNA in high-fat-fed mice resulted in stepwise reductions in both apoB and triglyceride secretion. Mipomersen lowered all apoB lipoproteins without reducing the production rate of either VLDL apoB or triglyceride. Our human data are consistent with long-standing models of posttranscriptional and posttranslational regulation of apoB secretion and are supported by *in vitro* and *in vivo* experiments. Targeting apoB synthesis may lower levels of apoB lipoproteins without necessarily reducing VLDL secretion, thereby lowering the risk of steatosis associated with this therapeutic strategy.

INTRODUCTION

Dyslipidemia, a major risk factor for cardiovascular disease (CVD), is characterized by elevated levels of apolipoprotein B100 (apoB) lipoproteins, including very-low-density lipoproteins (VLDL), carrying both triglycerides (TGs) and cholesterol, and low-density lipoproteins (LDL) carrying cholesterol (1). Although there is some heterogeneity in published results, increased secretion of apolipoprotein B (apoB) lipoproteins, particularly VLDL, is the characteristic abnormality observed in people with dyslipidemia (2, 3). On the basis of numerous clinical trials, however, lowering LDL cholesterol (LDL-C) remains the first-line therapy for reducing risk of CVD in such individuals (4). HMG-CoA (3-hydroxy-3-methylglutaryl coenzyme A) reductase inhibitors, better known as statins, are the most potent drugs available for reducing levels of apoB lipoproteins, mainly LDL, but also, to a lesser degree, VLDL. Although some studies have shown that statins can reduce production rates (PRs) of VLDL and LDL apoB, the central actions of statins result in an increase in the number of LDL receptors (LDLR) at the plasma membranes of cells, particularly the liver (5). More than 10% of individuals receiving statins are, however, clinically intolerant to these agents or can only take low doses of statin because of drug-specific side effects (6). Thus, about 50% of the patients on maximally tolerated statin therapy do not reach the recommended LDL-C levels established by National Cholesterol Education Program Adult Treat-

ment Panel III guidelines, especially patients with genetic lipid disorders such as familial hypercholesterolemia (7). Interest remains high, therefore, in the development of other therapeutic approaches to reduce circulating levels of apoB lipoproteins.

Two such agents—one a small-molecule inhibitor of microsomal triglyceride transfer protein (MTP) (8) and the other a second-generation antisense oligonucleotide (ASO) to apoB (9)—were recently approved by the U.S. Food and Drug Administration (FDA) for patients with homozygous familial hypercholesterolemia. Despite the ability of both drugs to reduce apoB lipoproteins, there are concerns about the occurrence of hepatic steatosis. Preclinical studies in rodents with either an ASO against MTP or small-molecule MTP inhibitors resulted in significant increases in liver TG levels (10, 11). This adverse effect was confirmed in studies of homozygous familial hypercholesterolemia patients with the recently approved MTP inhibitor, lomitapide (JUXTAPID, Aegerion) (8, 12). In preclinical studies in mice treated with ASO to apoB, there was no hepatic steatosis (10, 13), although increased liver TG has been observed in clinical trials of patients receiving mipomersen (KYNAMRO, Sanofi-Genzyme)—a fully phosphorothioate 20mer oligonucleotide with 5 2'-methoxyethyl residues at the 5' and 3' poles and a 10 deoxynucleotide center—for as long as 26 weeks for the treatment of familial hypercholesterolemia (14, 15). A combined analysis of three randomized trials with mipomersen treatment of patients with familial hypercholesterolemia indicated stabilization of steatosis during long-term treatment of more than 2 years. Reversal to baseline levels of hepatic fat was demonstrated in a subset of about 25% of participants who had magnetic resonance imaging performed 24 weeks after cessation of treatment (16).

The unique mechanisms of lomitapide or mipomersen to inhibit apoB production could explain the marked differences in the accumulation

¹Columbia University College of Physicians and Surgeons, 630 West 168th Street, New York, NY 10032, USA. ²Biological Sciences Department, New York City College of Technology, 300 Jay Street, Brooklyn, NY 11201, USA. ³Sanofi-Genzyme, 500 Kendall Street, Cambridge, MA 02142, USA.

*Corresponding author. E-mail: gr2104@cumc.columbia.edu (G.R.-S.); hng1@cumc.columbia.edu (H.N.G.)

†Present address: Ironwood Pharmaceuticals, 301 Binney Street, Cambridge, MA 02142, USA.

RESEARCH ARTICLE

of hepatic TG that have been observed in mice. When lomitapide inhibits MTP, the absence of transfer of TG, as well as phospholipids and cholesteryl esters, from the endoplasmic reticulum (ER) membrane to nascent apoB leads to degradation of apoB and the absence of lipid droplets in the ER lumen (17). In contrast, when apoB synthesis is submaximally inhibited, MTP is still able to transfer ER membrane lipids to apoB that escaped inhibition and was synthesized. This would be the case with the 200 mg/week dose of mipomersen that has been approved for patients with homozygous familial hypercholesterolemia as an adjunct to lipid-lowering medications and diet (18, 19). With the current dose of mipomersen used clinically, there could be increased “loading” of TG onto each apoB that is synthesized, resulting in the assembly and secretion of larger VLDL carrying more TG per particle.

Evidence supporting the ability of the liver to vary the amount of TG incorporated into nascent VLDL derives from studies of mice with increased de novo lipogenesis (20) and humans on high-carbohydrate diets (21) or with hepatic steatosis (22). This mechanism, which could compensate for reduced synthesis of apoB, would be unlikely to occur with inhibition of MTP, which is absolutely essential for the transfer of TG onto nascent apoB. An alternative or additional compensatory response to mipomersen-mediated decreases in the synthesis of apoB is suggested by many studies in hepatocytes demonstrating that only a portion of newly synthesized apoB is secreted, with the availability of hepatic lipids, particularly TG, being the major determinant of whether nascent apoB is secreted or degraded (17). In this model, submaximal inhibition of apoB synthesis might be compensated for by increasing the proportion of newly synthesized apoB that is secreted, thereby maintaining hepatic lipid homeostasis.

Here, we examined the effects of mipomersen treatment on fractional catabolic rates (FCRs) and PRs of apoB-containing lipoproteins and VLDL TG, as well as on hepatic de novo lipogenesis, in healthy volunteers. We also conducted studies of apoB and TG secretion during inhibition of apoB synthesis in a human hepatoma line and in mice. Our hypothesis was that mipomersen would decrease VLDL apoB PR, whereas VLDL TG PR would be unchanged, indicating the assembly and secretion of larger, TG-enriched VLDL particles. If our hypothesis is correct, mipomersen therapy could lower the levels of apoB-containing lipoproteins without necessarily causing significant hepatic steatosis.

RESULTS

Effects of mipomersen on lipid and apoB levels

Thirty-one subjects were screened, 20 enrolled, and 8 healthy men and 9 healthy women completed the study (table S1). Baseline characteristics are in Table 1. Participants were not taking medications known to affect lipid metabolism and had normal plasma glucose levels and normal hepatic and renal function. They received three weekly doses of placebo followed by seven weekly doses of mipomersen treatment (fig. S1). Mipomersen treatment resulted in significant reductions in plasma levels of total cholesterol, TG, LDL-C, and apoB, but not high-density lipoprotein cholesterol (HDL-C) (Table 2). The cholesterol and TG concentrations in VLDL, IDL, and LDL were decreased significantly by mipomersen treatment (Fig. 1, A and B), and there were marked reductions in the levels of VLDL apoB (–29%), IDL apoB (–25%), and LDL apoB (–42%) (Fig. 1C). We observed reductions in VLDL, IDL, and LDL particle numbers as determined by ion mobility (23) (table S2).

Table 1. Subject baseline characteristics. Twenty individuals were enrolled and 17 completed the study (table S1).

Mean age, years (±SD)	43.5 (14.2)
Gender, n (%)	
Male	8 (47.1)
Female	9 (52.9)
Race (%)	
White	29.4
Black	47.1
Asian	5.9
Unknown/mixed race	17.6
Mean BMI (±SD)	27.6 (3.2)

Table 2. Plasma lipid levels after 3 weeks of placebo and 7 weeks of mipomersen treatment. Data are means (±SD) of one sample obtained from each subject (*n* = 17) just before the kinetic study in each period. Percent change is based on the difference between levels during mipomersen treatment and during placebo treatment for each subject. The significance of percent change for each endpoint was examined by paired *t* test after determining that the data were normally distributed using the Kolmogorov-Smirnov normality test.

Parameter	Placebo (mg/dl)	Mipomersen (mg/dl)	Change (%)	<i>P</i> value
Total cholesterol	187.2 (25.7)	132.4 (29.7)	–28.6 (16.1)	<0.001
TG	86.9 (49.3)	57.1 (30.4)	–28.0 (28.0)	<0.001
LDL-C	114.7 (22.3)	61.9 (24)	–45.5 (18.9)	<0.001
ApoB	69.6 (18.9)	40.8 (14.1)	–40.3 (17.5)	<0.001
HDL-C	55.2 (14.3)	59.1 (17.9)	6.8 (19.3)	0.17

Effects of mipomersen on apoB metabolism in VLDL, IDL, and LDL

The reduction in VLDL apoB concentration during mipomersen treatment compared to placebo (Fig. 1C) was due to a 23% increase in the median FCR of VLDL apoB (Table 3). There was no significant effect of mipomersen treatment on VLDL apoB PR. The percent of VLDL apoB converted to IDL declined from 70 to 58%. This change, which reflects direct removal of VLDL apoB from plasma during mipomersen treatment, was associated with a significant 33% reduction in IDL apoB PR (Table 3) associated with the significant reduction in IDL apoB level (Fig. 1C) on mipomersen.

The FCR of IDL apoB did not change, and there was no effect of mipomersen on the conversion of IDL apoB to LDL (76 and 74%, respectively) (Table 3). However, the reduced IDL apoB PR, together with a modest fall in the secretion of LDL-like particles directly from the liver, resulted in a significant fall in LDL apoB concentration (Fig. 1C) owing to a 27% reduction in mean LDL apoB PR on mipomersen (Table 3). There was no significant change in total apoB PR (sum of VLDL apoB PR and direct LDL apoB PR) on mipomersen (Table 3).

The marked fall in LDL apoB levels during mipomersen therapy (Fig. 1C) was due not only to decreased PR but unexpectedly to a 30% increase in the FCR of LDL apoB (Table 3). Because circulating

Fig. 1. Effects of mipomersen treatment on human lipoprotein lipids and apoB levels. Blood samples were obtained from subjects during their kinetic studies performed after 3 weeks of treatment with placebo and after 7 weeks of treatment with mipomersen. Plasma was separated from those samples, and VLDL, IDL, and LDL were isolated by sequential ultracentrifugation. (A and B) Cholesterol and TG were measured by enzymatic methods. (C) ApoB was determined by commercial enzyme-linked immunosorbent assay. Data are means $\pm$ SD ($n = 5$ samples obtained from each of the 17 subjects during their two kinetic studies). P values determined by paired t tests after determining that the data were normally distributed using the Kolmogorov-Smirnov normality test.

proprotein convertase subtilisin/kexin type 9 (PCSK9) is an important regulator of hepatic LDLR (24), we quantified this protein in plasma. There was no significant effect of mipomersen on circulating concentrations of PCSK9 (261.7 ± 63.6 ng/ml on placebo versus 241.2 ± 58.2 ng/ml on mipomersen) ($P = 0.57$; paired t test). Alterations in LDL particle size, particularly leading to a greater proportion of medium-size particles, could increase affinity for the LDL receptor (25). However, mipomersen treatment was associated with a shift in the distribution of LDL toward small particles (table S3). Conversely, reductions in the numbers of all larger-size LDL subfractions (table S4) suggested increased numbers of LDL receptors during administration of mipomersen, with preferential removal of larger particles. Changes in hepatic cholesterol balances could alter LDL receptor levels: mipomersen had no effects on plasma levels of lathosterol, a marker of cholesterol synthesis, or either campesterol or β -sitosterol, both markers of cholesterol absorption (table S5).

Changes in FCRs of VLDL and LDL could have been affected by mipomersen-mediated changes in apoC-III or apoE concentrations.

There were no changes in plasma apoE levels from placebo (4.9 ± 1.5 mg/dl) to mipomersen treatment (4.5 ± 1.3 mg/dl) ($P = 0.2$; paired t test). In contrast, there was a fall in plasma apoC-III of 20% from placebo (9.3 ± 2.7) to mipomersen (7.3 ± 3.2) ($P = 0.02$; paired t test).

Effects of mipomersen on VLDL TG metabolism

The concentration of VLDL TG fell by 20% during mipomersen treatment (Fig. 1B). This was accompanied by a 46% increase in the mean FCR of VLDL TG during mipomersen treatment, with no change seen in the PR of VLDL TG (Table 3). The percentage of VLDL TG produced from hepatic de novo lipogenesis did not change ($11.2 \pm 6.3\%$ and $10.5 \pm 5.6\%$ during placebo and mipomersen, respectively) ($P = 0.50$; paired t test). To gain insight into the increase in VLDL TG FCR, we measured post-heparin hepatic lipase (HL) and lipoprotein lipase (LpL) activities in plasma. Neither of these enzymes was affected by mipomersen treatment (table S6). Fasting plasma levels of fatty acid (FA) and β -hydroxybutyrate also did not change between periods (table S6). The individual FCRs and PRs for VLDL, IDL, and LDL apoB and VLDL TG, for the 17 subjects, are in table S7.

We hypothesized that mipomersen treatment might be associated with a shift in the size distribution of secreted VLDL, with a larger TG-rich VLDL entering the circulation. The ratio of PRs of VLDL TG to VLDL apoB, which should reflect the size of newly secreted VLDL particles, increased by a median value of 17%, indicating assembly and secretion of larger VLDL (Table 3). These data were supported by a significant increase of 25% in the ratio of TG/apoB in circulating VLDL during mipomersen treatment (Table 3).

Effects of siRNA knockdown of APOB in HepG2 cells on apoB secretion

To better understand the lack of effect of mipomersen on the secretion of VLDL apoB in vivo, we examined the effects of the knockdown of APOB on the secretion of apoB in the human hepatoma cell line HepG2 in vitro. For these studies, we used a small interfering RNA (siRNA) that has been extensively validated against human APOB in vitro in HepG2 cells (26) and in vivo in nonhuman primates (27) to titrate the knockdown of APOB mRNA. We determined that maximal siRNA-mediated knockdown of APOB mRNA was attained at a concentration of 10 nM, with about 80% inhibition of apoB synthesis (cell) and 60% inhibition of apoB secretion (media) during a 2-hour steady-state labeling study compared to a scrambled siRNA control (Fig. 2A). Titration of the siRNA between 0.05 and 1.0 nM resulted in dose-related reductions of APOB mRNA (Fig. 2B) and synthesis of the protein in short-pulse and pulse-chase experiments with or without inhibition of the proteasome (Fig. 2C). There was an obvious relationship between increasing siRNA-mediated knockdown of APOB expression and cellular accumulation of newly synthesized apoB in 2-hour labeling studies (Fig. 2D). However, secretion of apoB remained constant across the doses of siRNA (Fig. 2E).

When we depicted secretion of apoB as the percent of the total newly synthesized apoB in cells plus the media after 2 hours of radio-labeling, there was increased efficiency of secretion (Fig. 2F). In contrast, a small-molecule inhibitor of MTP caused parallel decreases in cellular (Fig. 2G) and secreted (Fig. 2H) apoB after 2 hours of radio-labeling. Thus, with MTP inhibition, there was no change in the efficiency of secretion of apoB with increasing inhibition of MTP (Fig. 2I). siRNA knockdown of APOB did not affect expression of LDL

RESEARCH ARTICLE

Table 3. ApoB kinetics. The effects of mipomersen on apoB metabolism were determined from 17 kinetic studies at the end of the placebo and mipomersen treatment periods. Absolute values for the placebo and mipomersen periods are shown. If the data were found to be normally distributed by the Kolmogorov-Smirnov test, they are presented as means ($\pm$ SD),

and statistical analyses were performed on percent change for each endpoint on mipomersen from placebo using paired *t* tests. If the data were not normally distributed, the median and interquartile values (in parentheses) are shown for percent change, and the Wilcoxon rank-sum test was performed to test for significance.

Parameter	Placebo	Mipomersen	Change (%)	P value
VLDL apoB FCR (pools/day)	7.6 (3.6)	9.9 (4.6)	23.0 (-7.4, 55.6)	0.023
VLDL apoB PR (mg kg ⁻¹ day ⁻¹)	21.7 (11.3)	19.0 (10.9)	-17.6 (-32.7, -0.9)	0.15
VLDL to IDL (% conversion)*	70.2 (24.2)	58.2 (24.5)	-12.1	0.09
IDL apoB FCR (pools/day)	5.5 (3.5)	5.8 (3.6)	11.1	0.30
IDL apoB PR (mg kg ⁻¹ day ⁻¹)	15.0 (8.1)	10.2 (6.9)	-32.7 (-53.5, -5.7)	0.05
IDL to LDL (% conversion)*	76.2 (23)	74.4 (26)	-1.7	0.84
LDL apoB FCR (pools/day)	0.52 (0.14)	0.65 (0.19)	29.7	<0.001
LDL apoB PR (mg kg ⁻¹ day ⁻¹)	12.7 (4.6)	8.8 (3.1)	-26.5	0.001
Direct LDL PR (mg kg ⁻¹ day ⁻¹) [†]	2.6 (3.5)	2.2 (1.5)	0.0 (-0.8, 1.5)	0.89
LDL apoB PR from direct LDL secretion (%)*	21.3 (22)	26.8 (18)	5.5	0.38
Total apoB PR (sum VLDL PR and direct LDL PR) (mg kg ⁻¹ day ⁻¹)	24.3 (11.5)	21.3 (10.7)	-15.1 (-34.5, -7.4)	0.26
VLDL TG FCR (pools/hour)	10.5 (7.9)	13.2 (8.4)	46.3	0.03
VLDL TG PR (mg kg ⁻¹ hour ⁻¹)	13.1 (6.4)	13.3 (7.0)	10.0	0.40
VLDL TG PR/VLDL apoB PR ratio	22.9 (27.0)	26.2 (37.0)	16.5 (-7.8, 93.2)	0.05
Plasma VLDL TG/VLDL apoB ratio	14.5 (13.7)	15.3 (9.0)	24.6	0.03

*For the percent conversions (VLDL to IDL; IDL to LDL), absolute differences between placebo and mipomersen results were used to test for significance.

[†]Some direct LDL secretion rates were zero, so no percent change was calculated for this parameter; instead, values are absolute differences.

receptor (*LDLR*), inducible degrader of the *LDLR* (*IDOL*), or *PCSK9* in HepG2 cells (fig. S2).

These results demonstrate that HepG2 cells, under culture conditions where there is very little intracellular TG, are able to accommodate submaximal reductions in apoB synthesis by increasing the proportion of newly synthesized apoB that is secreted.

Effect of ASO knockdown of *ApoB* in mice on secretion of apoB and TG

To further support the validity of our observation of a lack of effect of mipomersen on VLDL apoB secretion in normal humans, we performed similar experiments in vivo in *ApoBec-1* knockout mice, using a murine-specific *ApoB* ASO. *ApoBec-1* knockout mice were used because they synthesize and secrete only apoB from their livers, making them more like humans. Mice on either chow or a high-fat diet (HFD) for 6 weeks were treated with *ApoB* ASO in doses ranging from 10 to 50 mg/week for an additional 3 weeks, while continuing their diets, at which time apoB and TG secretion were measured. The doses were chosen from preliminary studies where 200 or 100 mg/week of *ApoB* ASO significantly reduced both TG and apoB secretion in chow-fed mice (fig. S3).

In the titration experiment, *ApoB* ASO reduced hepatic *ApoB* mRNA levels in both groups of mice in a dose-dependent manner, with the

response more pronounced in the HFD-fed mice (Fig. 3A), similar to the results of Lee *et al.*, who conducted experiments with the same ASO in mice on chow or HFD (10). Relevant to this experiment is the evidence that *ApoB* ASO does not significantly alter *ApoB* mRNA levels after 2 weeks of treatment (13) in the small intestine, and although it does lower *ApoB* mRNA after 6 to 8 weeks of treatment, APOB protein is not altered at those times (10, 13). The effects of ASO treatment on apoB and TG secretion demonstrated different patterns in chow- versus HFD-fed mice. Whereas *apoB* secretion was unaffected by *apoB* ASO at the doses administered in the chow-fed mice, it was inhibited in a stepwise, dose-dependent manner in the HFD mice (Fig. 3B). Furthermore, *apoB* ASO had modest, non-dose-related effects on TG secretion in the chow-fed mice, but caused greater reductions of TG secretion in the HFD mice with a clear trend for dose responsiveness (Fig. 3C). The hepatic TG concentration in chow-fed mice receiving control ASO was 159 ± 33 μ g/mg protein and, as expected, much higher (625 ± 207 μ g/mg protein) in the HFD group on control ASO ($P < 0.001$, by unpaired *t* test).

Hepatic steatosis in the HFD group was associated with greater absolute rates of secretion of newly synthesized apoB in the control ASO-treated mice compared to chow-fed control ASO-treated mice (fig. S4). *ApoB* ASO treatment did not affect expression of the genes for *Ldlr*, *Idol*, or *Pcsk9* (fig. S5). Furthermore, *ApoB* ASO treatment

RESEARCH ARTICLE

Fig. 2. Effects of siRNA-mediated knockdown of *APOB* on apoB secretion in human HepG2 cells. (A) *APOB* siRNA (10 nM) inhibits the synthesis and secretion of apoB in HepG2 cells. HepG2 cells were transfected with 10 nM *APOB* or 10 nM irrelevant control (Con) siRNA for 2 days, radiolabeled continuously with [35 S]methionine/cysteine for 2 hours, and radioactivity [counts per minute (CPM)] in cellular or media apoB was determined. (B) *APOB* siRNA knocks down *APOB* mRNA in a dose-dependent manner. HepG2 cells were transfected at varying doses of *APOB* siRNA for 2 days. Total RNA was extracted and *APOB* mRNA levels were analyzed by quantitative polymerase chain reaction (qPCR). (C) *APOB* siRNA reduces initial synthesis rates of apoB in a dose-dependent manner. Cells were labeled with [35 S]methionine/cysteine for 10 min and chased for 10 min with or without proteasome inhibitor ALLN pretreatment for 1 hour. (D and E) The effect of *APOB* siRNA doses on cellular and secreted apoB in HepG2

cells transfected as in (B), and apoB synthesis and secretion analyzed as in (A). (F) Efficiency of apoB secretion as a function of *APOB* siRNA knockdown. Efficiency was defined as the percentage of apoB secreted into the medium compared with the amount of total newly synthesized apoB (cellular and secreted apoB over 2 hours). (G and H) Cellular and secreted apoB in HepG2 cell culture after treatment with an MTP inhibitor (MTPi) for 1 hour (before steady-state labeling for 2 hours). (I) Efficiency of apoB secretion as a function of MTP inhibition. Data were plotted using the cell and secreted apoB data, as described in (F). All data are mean percentage of control [siRNA (1 nM) in (B) to (F); dimethyl sulfoxide (DMSO) in (G) to (I)] $\pm$ SD ($n = 3$). * $P < 0.05$; ** $P < 0.01$ versus respective control, unless otherwise noted, one-way analysis of variance (ANOVA) followed by post hoc comparisons using the Fisher LSD (least significant difference) method.

did not alter the level of LDLR protein in the livers of mice on either diet (fig. S6).

These in vivo data are consistent with greater need to use a larger proportion of newly synthesized apoB for assembly with TG to secrete VLDL when there is steatosis. In this situation, therefore, any reduction of apoB synthesis will result in decreased apoB secretion. Indeed, an exploratory analysis of the relationship between baseline VLDL apoB production and the decrease in VLDL apoB on mipomersen in the human study indicated just such a relationship, although the correlation did not achieve significance (fig. S7).

DISCUSSION

Mipomersen is a second-generation ApoB ASO that inhibits the synthesis of apoB (13, 28). In a phase 1 study, treatment with 50 to 400 mg of mipomersen injected subcutaneously weekly produced dose-dependent reductions in plasma LDL-C and apoB concentrations (29). On the basis of those results and phase 2 studies (18, 19), a dose of 200 mg per week, by subcutaneous injection, was chosen for further development and was the dose approved by the FDA for treatment of patients with homozygous familial hypercholesterolemia (9). The phase 1 and 2 data

Fig. 3. Effects of ApoB ASO treatment on apoB and TG secretion in *Apobec-1* knockout mice. Chow- and HFD-fed mice were treated for 3 weeks with ApoB ASO. (A) ApoB mRNA expression was determined using qPCR of livers from both chow and HFD mice and normalized to liver β -actin mRNA. In each diet group, irrelevant, control (Ctrl) ASO-treated mRNA levels were set as 100%. (B) Mice were fasted 4 hours and injected with Tyloxapol and [35 S]methionine intravenously. Plasma [35 S]-labeled apoB levels in CPM were determined in samples obtained at 120 min after the start of the study. ApoB radioactivity in mice receiving the irrelevant control ASO was set as 100%. (C) Blood samples were collected at 0, 30, 60, 90, and 120 min for measurement of TG levels. TG secretion rate was determined by the change in plasma TG concentration between 30 and 120 min after Tyloxapol injection and divided by 1.5. The secretion rate of the mice receiving irrelevant control ASO was set at 100%. Data in (A) to (C) are means $\pm$ SD (n noted on figure). * $P < 0.05$ versus control, unless otherwise noted, one-way ANOVA and Tukey's post hoc test.

indicated that the 300- and 400-mg weekly doses had greater LDL-C and apoB-lowering efficacy than the 200-mg weekly dose, which was the dose used here. Thus, it is reasonable to conclude that the 200-mg weekly dose results in submaximal reductions in apoB synthesis; this, we believe, is critical to the interpretation of our results.

As expected, mipomersen significantly reduced plasma levels of LDL-C and apoB. VLDL, IDL, and LDL apoB concentrations were also reduced, as were levels of VLDL TG. The reductions in LDL apoB levels resulted from decreased LDL PRs, but unexpectedly, the fall in LDL PR did not result from decreased VLDL apoB secretion into plasma, which was our primary hypothesis. Instead, decreased production of LDL resulted from increased direct clearance of VLDL apoB with less conversion to IDL, the immediate precursor of LDL. A modest reduction

in the direct secretion of LDL into plasma during mipomersen treatment also contributed to the fall in LDL PR. Additionally, and also unexpectedly, an increase in FCR contributed to the reduced LDL apoB levels present during mipomersen treatment.

Why would treatment with a drug that reduces *APOB* mRNA, thereby reducing apoB synthesis, not lower VLDL apoB PR? In vitro studies in a variety of primary hepatocytes and hepatic cell lines have demonstrated repeatedly that apoB is typically synthesized constitutively and that its assembly into a VLDL and secretion involves the co-translational lipidation of the nascent polypeptide as it traverses the ER (17). The latter step plays a crucial role in determining how much VLDL is secreted; in the absence of adequate lipids or MTP activity, co-translational and co-translocational degradation of apoB results (17). Additional sites of intracellular degradation of apoB can also occur during or after the assembly of apoB with lipids (30), adding further complexity to the secretory process. The extensive post-transcriptional and posttranslational control of the assembly and secretion of VLDL results in a range of secretion from 20 to 70% of newly synthesized apoB from cultured hepatocytes. The relevance of these cell-based studies to human physiology is supported by the wide range of secretory rates of VLDL in normal and hyperlipidemic individuals, and the ability to alter VLDL PR in individuals with different diets and drugs (31, 32).

With these data in mind, we carried out a series of studies in HepG2 cells in which we reduced *APOB* mRNA in a stepwise fashion with siRNA (26, 27). Our findings of increased efficiency of apoB secretion across a range of inhibition of apoB synthesis strongly support our in vivo results. For example, if some-

one with normal liver TG levels makes 100 apoB molecules per minute and only needs to secrete 50 VLDL particles (with one apoB per particle) to maintain hepatic lipid homeostasis, inhibition of synthesis by 40%, to 60 molecules of apoB produced per minute, should allow the liver to maintain hepatic lipids at pretreatment levels by increasing the proportion of nascent apoB that is secreted. In contrast, if hepatic lipids are elevated and a greater percentage of newly synthesized apoB is constantly being secreted as the liver attempts to reduce hepatic TG, inhibiting synthesis of apoB would likely lead to decreased secretion of VLDL.

This is what we observed in HFD-fed mice compared to chow-fed mice. The HFD mice had steatosis and much higher rates of apoB secretion when treated with control ASO and had dose-dependent reductions in apoB secretion when treated with apoB ASO; this was in

marked contrast to what was observed in chow-fed mice. Results of an exploratory analysis in our subjects provided support for our model of the role of apoB in maintaining hepatic lipid homeostasis. In our normolipidemic participants, higher VLDL apoB PRs on placebo were associated with greater reductions in VLDL apoB secretion on mipomersen. Further studies of the effects of mipomersen on apoB metabolism in people with hepatic steatosis and a dyslipidemia characterized by increased VLDL apoB and VLDL TG secretion will be required to definitively address this important question.

Our second hypothesis, that VLDL TG PR would not be affected by mipomersen owing to assembly and secretion of larger VLDL, was supported by our findings of increased ratios of TG to apoB in circulating VLDL and an increased ratio of VLDL TG PR to VLDL apoB PR. A rigorous examination of this hypothesis, which is supported by in vitro studies in cultured liver cells that were treated with FAs (33) and in vivo studies in mice and humans in which hepatic de novo lipogenesis was increased (20, 34), must await further studies of mipomersen treatment of hypertriglyceridemic subjects. Plasma and VLDL TG levels fell despite any change in VLDL TG PR. Our finding of increased FCRs for VLDL TG in the absence of increases in post-heparin LpL and HL activities points to previous studies showing increased efficiency of LpL-mediated lipolysis of larger and more TG-enriched particles (35). Additionally, we observed an increase in both the FCR of VLDL apoB and the direct removal of VLDL apoB without conversion to IDL or LDL. The increased FCR of VLDL apoB could simply have been the result of a more efficient LpL-mediated lipolysis of larger VLDL TG or a result of reduced levels of apoC-III we observed during mipomersen treatment. However, that would have resulted in a more rapid and efficient conversion of VLDL to IDL and LDL, rather than the decreased conversion we observed.

Both apoE and apoC-III affect hepatic removal of VLDL; apoE levels in plasma were not altered by mipomersen treatment, but the reductions in apoC-III levels we observed could have resulted in greater removal of VLDL or their remnants by the liver with less conversion to IDL. The latter would be more likely if the reduced apoC-III levels resulted from mipomersen-mediated decreased secretion of the protein by the liver rather than those lower concentrations being the result of lower levels of VLDL, IDL, and LDL, which all carry apoC-III. Further studies will be required to determine if decreased apoB synthesis affects apoC-III synthesis or secretion by the liver.

The reductions in LDL we observed resulted from both decreased PR due to less conversion of VLDL to IDL and LDL, and a significant and unexpected increase in LDL FCR. We were unable to discern why inhibition of hepatic apoB synthesis would lead to an increase in LDL FCR, which is typically associated with increased levels of hepatic LDL receptors. However, we did measure plasma PCSK9 concentrations as a surrogate, because lower PCSK9 levels would mean increased hepatic LDL receptors available to bind LDL (24). Mipomersen had no effect on plasma PCSK9 levels in humans. Reduced rates of endogenous cholesterol synthesis or intestinal absorption of cholesterol could also lead downstream to increased LDLR, but we saw no effects of mipomersen treatment on the three markers of the whole-body cholesterol metabolism: lathosterol, campesterol, or β -sistosterol. *LDLR*, *PCSK9*, and *IDOL* mRNA were unchanged in human HepG2 cells treated with maximally suppressive doses of siRNA. Furthermore, mipomersen did not affect mRNA levels of the *Ldlr*, *Pcsk9*, and *Idol*, or the level of LDLR protein in mice. An explanation that is not dependent on the number of LDL receptors in the liver would be greater affinity of LDL for the same

number of receptors; we did see a change in the distribution of LDL size, in the direction of smaller LDL particles, but small LDLs have lower affinity for the LDLR. The reductions that occurred in the numbers of particles in all the large LDL subfractions are consistent, however, with increased LDLR-mediated clearance. Additionally, the cholesterol-to-apoB ratio in LDL increased; larger, cholesterol-enriched LDL is believed to have greater affinity for the LDLR. As noted above, we observed reductions in apoC-III, which can inhibit the uptake of LDL by its receptor. However, apoC-III is only carried on a small fraction of LDL, and the changes we observed in apoC-III are unlikely to explain the significant increase in LDL FCR with mipomersen treatment. Overall, we do not have a clear explanation for the increased LDL apoB FCR during mipomersen treatment.

We understand that the results of mipomersen treatment of humans with homozygous familial hypercholesterolemia suggest that the LDL receptor is not needed for efficacy of this agent (9). Studies of apoB metabolism in homozygous familial hypercholesterolemia with null mutations have demonstrated increased secretion of apoB lipoproteins (36, 37); in our model, mipomersen inhibition of apoB synthesis would decrease apoB secretion in those individuals. In summary, our study of the effects of mipomersen in healthy subjects, supported by our studies in HepG2 cells and mice, provides new and unique insights into mechanisms for the maintenance of hepatic lipid homeostasis when apoB synthesis is reduced. Our results indicate that inhibition of apoB synthesis should remain a target for therapies to reduce levels of all atherogenic lipoproteins.

MATERIALS AND METHODS

Study design

This was a single blind, two-period, linear, placebo-controlled, phase 1 study in healthy volunteers. The first primary hypothesis was that mipomersen, administered at a dose of 200 mg weekly, would significantly reduce the PR of VLDL apoB. The second hypothesis was that there would be an increase in the efficiency of VLDL TG fractional clearance from the circulation during mipomersen treatment. Exploratory hypotheses included that there would not be a change in VLDL TG production; that there would not be a change in direct removal of VLDL apoB from the circulation; that LDL apoB production would be reduced, but fractional removal would not be affected; that hepatic DNL would not be affected; and that FA oxidation in the liver, as estimated by measurement of circulating β -hydroxybutyrate, would not change during mipomersen treatment. The primary endpoint was the percent change in VLDL apoB PR from the end of placebo to the end of mipomersen treatment. The sample size calculation indicated that 16 subjects would provide 80% power to detect a 20% reduction in VLDL apoB PR, on the basis of a paired *t* test with a 5% type I error rate, assuming an SD of about 6 mg kg⁻¹ day⁻¹ as observed in previous research (38, 39) and an expected VLDL apoB PR of about 25 mg kg⁻¹ day⁻¹ at the end of the placebo period. Additional details of the study protocol are in the Supplementary Materials and fig. S1.

Human volunteers

All subjects signed informed consent. For screening, see the Supplementary Materials. Subjects were nonsmokers, were of ages 18 to 75 years, had a body mass index (BMI) of less than 38 kg/m², and were not on any lipid-altering medications. All participants had blood glucose

RESEARCH ARTICLE

<115 mg/dl, TG <170 mg/dl, and LDL-C >100 mg/dl and <160 mg/dl if they had more than two cardiovascular risk factor. Subjects who met the eligibility criteria were instructed to follow an American Heart Association–recommended diet. After an additional week, subjects returned to the Irving Institute for Clinical and Translational Research (IICTR) for study randomization and fasting measurements of plasma lipids, a chemistry panel, hematology studies, and vital signs.

Mathematical modeling of the apoB and TG enrichment data

ApoB kinetics in VLDL, IDL, and LDL were estimated using the leucine and phenylalanine enrichment data from the constant infusion and the bolus injection, respectively (38, 39). The kinetics of VLDL TG were analyzed using the enrichment data generated by the injection of D5-glycerol. Mathematical modeling is described in the Supplementary Methods.

In vitro studies of APOB knockdown and inhibition of MTP activity in HepG2 cells

HepG2 cells were obtained from American Type Culture Collection and cultured in a growth medium of Dulbecco's modified Eagle's medium (high glucose, 5% antibiotics, and 10% fetal bovine serum). Transfection of HepG2 cells and metabolic labeling to measure the secretion and synthesis of apoB were performed with both steady-state and pulse-chase labeling protocols as described previously in our laboratory; details are provided in the Supplementary Materials (40, 41).

In vivo studies of ApoB knockdown in mice

All animal experiments were approved by the Columbia University Institutional Animal Care and Use Committee. Details regarding the mice are in the Supplementary Materials. After 6 weeks on HFD, *ApoBec-1* knockout mice were started on treatment with mouse control ASO (ISIS #141923) or a second-generation mouse apoB ASO (Ionis Pharmaceuticals #147764) for 3 weeks; these reagents were provided by Ionis Pharmaceuticals. Mice remaining on chow diet were started on ASO treatment at the same age as the mice on HFD. The *ApoB* ASO was administered at different doses (10, 25, or 50 mg/week) mixed with control ASO to give equal amounts of ASO in each injection. At the end of 3 weeks of ASO treatment, mice were fasted for 4 hours, and TG and apoB secretion was measured by intravenously injecting Tyloxapol (500 mg/kg) (T8761-50G, Sigma-Aldrich) and 200 μ Ci per mouse [35 S]methionine (1175 Ci/mmol; PerkinElmer Life Sciences) as described previously (42).

Statistical analysis

The Wilcoxon signed-rank test and the paired *t* test were used to test the hypothesis that there was no change between the first and second turnover studies. The Kolmogorov-Smirnov statistic tested the hypothesis that the percent/nominal change was normally distributed. For the in vitro cell and in vivo mouse studies, all data are presented as means $\pm$ SD. Differences in the means among multiple groups were compared for significance using a one-way ANOVA followed by individual comparisons when appropriate. *P* < 0.05 was considered significant.

SUPPLEMENTARY MATERIALS

www.sciencetranslationalmedicine.org/cgi/content/full/8/323/323ra12/DC1

Methods

Fig. S1. Study flow diagram.

Fig. S2. No changes in *LDLR*, *IDOL*, or *PCSK9* mRNA levels in HepG2 cells after knockdown of *APOB*.

Fig. S3. *ApoB* ASO inhibits hepatic secretion of TG and apoB from mice on either chow or HFD diet.

Fig. S4. ApoB secretion is increased in *ApoBec-1* knockout mice on HFD.

Fig. S5. *ApoB* siRNA treatment does not affect expression of *Ldlr*, *Idol*, or *Pcsk9* in mouse livers.

Fig. S6. Hepatic LDL receptor was not affected by *ApoB* ASO.

Fig. S7. Relationship between mipomersen-induced reductions in VLDL apoB PR and baseline VLDL PR.

Table S1. Participant information.

Table S2. Effect of mipomersen on apoB lipoprotein particle number by ion mobility analysis.

Table S3. Effect of mipomersen on apoB lipoprotein size distribution by ion mobility analysis.

Table S4. Effect of mipomersen on LDL size subfraction particle numbers by ion mobility analysis.

Table S5. Markers of cholesterol synthesis and absorption.

Table S6. Hepatic lipase, lipoprotein lipase, FFA, and β -hydroxybutyrate levels.

Table S7. Individual participant apoB kinetics of VLDL, IDL, LDL, and VLDL TG FCR.

References (43–49)

REFERENCES AND NOTES

- M. J. Chapman, H. N. Ginsberg, P. Amarenco, F. Andreotti, J. Borén, A. L. Catapano, O. S. Descamps, E. Fisher, P. T. Kovanen, J. A. Kuivenhoven, P. Lesnik, L. Masana, B. G. Nordestgaard, K. K. Ray, Z. Reiner, M.-R. Taskinen, L. Tokgözoğlu, A. Tybjaerg-Hansen, G. F. Watts; European Atherosclerosis Society Consensus Panel, Triglyceride-rich lipoproteins and high-density lipoprotein cholesterol in patients at high risk of cardiovascular disease: Evidence and guidance for management. *Eur. Heart J.* **32**, 1345–1361 (2011).
- M. Adiels, S.-O. Olofsson, M.-R. Taskinen, J. Borén, Overproduction of very low-density lipoproteins is the hallmark of the dyslipidemia in the metabolic syndrome. *Arterioscler. Thromb. Vasc. Biol.* **28**, 1225–1236 (2008).
- D. C. Chan, P. H. R. Barrett, G. F. Watts, Lipoprotein transport in the metabolic syndrome: Pathophysiological and interventional studies employing stable isotope and modelling methods. *Clin. Sci.* **107**, 233–249 (2004).
- Cholesterol Treatment Trialists' (CTT) Collaboration, C. Baigent, L. Blackwell, J. Emberson, L. E. Holland, C. Reith, N. Bhalra, R. Peto, E. H. Barnes, A. Keech, J. Simes, R. Collins, Efficacy and safety of more intensive lowering of LDL cholesterol: A meta-analysis of data from 170 000 participants in 26 randomised trials. *Lancet* **376**, 1670–1681 (2010).
- M. S. Brown, J. L. Goldstein, A receptor-mediated pathway for cholesterol homeostasis. *Science* **232**, 34–47 (1986).
- E. S. Stroes, P. D. Thompson, A. Corsini, G. D. Vladutiu, F. J. Raal, K. K. Ray, M. Roden, E. Stein, L. Tokgözoğlu, B. G. Nordestgaard, E. Bruckert, G. De Backer, R. M. Krauss, U. Laufs, R. D. Santos, R. A. Hegele, G. K. Hovingh, L. A. Leiter, F. Mach, W. März, C. B. Newman, O. Wiklund, T. A. Jacobson, A. L. Catapano, M. J. Chapman, H. N. Ginsberg; European Atherosclerosis Society Consensus Panel, Statin-associated muscle symptoms: Impact on statin therapy—European Atherosclerosis Society Consensus Panel Statement on Assessment, Aetiology and Management. *Eur. Heart J.* **36**, 1012–1022 (2015).
- A. K. Gitt, H. Drexel, J. Feely, J. Ferrières, J. R. Gonzalez-Juanatey, K. K. Thomsen, L. A. Leiter, P. Lundman, P. M. da Silva, T. Pedersen, D. Wood, C. Jünger, P. S. Dellea, V. Sazonov, F. Chazelle, J. J. P. Kastelein; DYSIS Investigators, Persistent lipid abnormalities in statin-treated patients and predictors of LDL-cholesterol goal achievement in clinical practice in Europe and Canada. *Eur. J. Prev. Cardiol.* **19**, 221–230 (2012).
- M. Cuchel, E. A. Meagher, H. du Toit Theron, D. J. Blom, A. D. Marais, R. A. Hegele, M. R. Averna, C. R. Sirtori, P. K. Shah, D. Gaudet, C. Stefanutti, G. B. Vigna, A. M. E. Du Plessis, K. J. Propert, W. J. Sasiela, L. T. Bloedon, D. J. Rader; Phase 3 HoFH Lomitapide Study investigators, Efficacy and safety of a microsomal triglyceride transfer protein inhibitor in patients with homozygous familial hypercholesterolaemia: A single-arm, open-label, phase 3 study. *Lancet* **381**, 40–46 (2013).
- F. J. Raal, R. D. Santos, D. J. Blom, A. D. Marais, M.-J. Chamg, W. C. Cromwell, R. H. Lachmann, D. Gaudet, J. L. Tan, S. Chasan-Taber, D. L. Tribble, J. D. Flaim, S. T. Crooke, Mipomersen, an apolipoprotein B synthesis inhibitor, for lowering of LDL cholesterol concentrations in patients with homozygous familial hypercholesterolaemia: A randomised, double-blind, placebo-controlled trial. *Lancet* **375**, 998–1006 (2010).
- R. G. Lee, W. Fu, M. J. Graham, A. E. Mullick, D. Sipe, D. Gattis, T. A. Bell, S. Booten, R. M. Crooke, Comparison of the pharmacological profiles of murine antisense oligonucleotides targeting apolipoprotein B and microsomal triglyceride transfer protein. *J. Lipid Res.* **54**, 602–614 (2013).
- A. S. Wierzbicki, T. Hardman, W. T. Prince, Future challenges for microsomal transport protein inhibitors. *Curr. Vasc. Pharmacol.* **7**, 277–286 (2009).
- M. Cuchel, L. T. Bloedon, P. O. Szapary, D. M. Kolansky, M. L. Wolfe, A. Sarkis, J. S. Millar, K. Ikewaki, E. S. Siegelman, R. E. Gregg, D. J. Rader, Inhibition of microsomal triglyceride transfer protein in familial hypercholesterolemia. *N. Engl. J. Med.* **356**, 148–156 (2007).

13. R. M. Crooke, M. J. Graham, K. M. Lemonidis, C. P. Whipple, S. Koo, R. J. Perera, An apolipoprotein B antisense oligonucleotide lowers LDL cholesterol in hyperlipidemic mice without causing hepatic steatosis. *J. Lipid Res.* **46**, 872–884 (2005).
14. M. E. Visser, F. Akdim, D. L. Tribble, A. J. Nederveen, T. J. Kwok, J. J. P. Kastelein, M. D. Trip, E. S. G. Stroes, Effect of apolipoprotein-B synthesis inhibition on liver triglyceride content in patients with familial hypercholesterolemia. *J. Lipid Res.* **51**, 1057–1062 (2010).
15. E. A. Stein, R. Dufour, C. Gagne, D. Gaudet, C. East, J. M. Donovan, W. Chin, D. L. Tribble, M. McGowan, Apolipoprotein B synthesis inhibition with mipomersen in heterozygous familial hypercholesterolemia: Results of a randomized, double-blind, placebo-controlled trial to assess efficacy and safety as add-on therapy in patients with coronary artery disease. *Circulation* **126**, 2283–2292 (2012).
16. R. D. Santos, P. B. Duell, C. East, J. R. Guyton, P. M. Moriarty, W. Chin, R. S. Mittleman, Long-term efficacy and safety of mipomersen in patients with familial hypercholesterolemia: 2-year interim results of an open-label extension. *Eur. Heart J.* **36**, 566–575 (2015).
17. H. N. Ginsberg, E. A. Fisher, The ever-expanding role of degradation in the regulation of apolipoprotein B metabolism. *J. Lipid Res.* **50**, S162–S166 (2009).
18. F. Akdim, E. S. G. Stroes, E. J. G. Sijbrands, D. L. Tribble, M. D. Trip, J. W. Jukema, J. D. Flaim, J. Su, R. Yu, B. F. Baker, M. K. Wedel, J. J. P. Kastelein, Efficacy and safety of mipomersen, an antisense inhibitor of apolipoprotein B, in hypercholesterolemic subjects receiving stable statin therapy. *J. Am. Coll. Cardiol.* **55**, 1611–1618 (2010).
19. F. Akdim, D. L. Tribble, J. D. Flaim, R. Yu, J. Su, R. S. Geary, B. F. Baker, R. Fuhr, M. K. Wedel, J. J. P. Kastelein, Efficacy of apolipoprotein B synthesis inhibition in subjects with mild-to-moderate hyperlipidaemia. *Eur. Heart J.* **32**, 2650–2659 (2011).
20. J. D. Horton, H. Shimano, R. L. Hamilton, M. S. Brown, J. L. Goldstein, Disruption of LDL receptor gene in transgenic SREBP-1a mice unmasks hyperlipidemia resulting from production of lipid-rich VLDL. *J. Clin. Invest.* **103**, 1067–1076 (1999).
21. J. Melish, N.-A. Le, H. Ginsberg, D. Steinberg, W. V. Brown, Dissociation of apoprotein B and triglyceride production in very-low-density lipoproteins. *Am. J. Physiol.* **239**, E354–E362 (1980).
22. M. Adiels, M.-R. Taskinen, C. Packard, M. J. Caslake, A. Soro-Paavonen, J. Westerbacka, S. Vehkavaara, A. Häkkinen, S.-O. Olofsson, H. Yki-Järvinen, J. Borén, Overproduction of large VLDL particles is driven by increased liver fat content in man. *Diabetologia* **49**, 755–765 (2006).
23. M. P. Caulfield, S. Li, G. Lee, P. J. Blanche, W. A. Salameh, W. H. Benner, R. E. Reitz, R. M. Krauss, Direct determination of lipoprotein particle sizes and concentrations by ion mobility analysis. *Clin. Chem.* **54**, 1307–1316 (2008).
24. N. G. Seidah, A. Prat, The biology and therapeutic targeting of the proprotein convertases. *Nat. Rev. Drug Discov.* **11**, 367–383 (2012).
25. H. Campos, K. S. Arnold, M. E. Balestra, T. L. Innerarity, R. M. Krauss, Differences in receptor binding of LDL subfractions. *Arterioscler. Thromb. Vasc. Biol.* **16**, 794–801 (1996).
26. M. Tadin-Strapps, L. B. Peterson, A.-M. Cumiskey, R. L. Rosa, V. H. Mendoza, J. Castro-Perez, O. Puig, L. Zhang, W. R. Strapps, S. Yendluri, L. Andrews, V. Pickering, J. Rice, L. Luo, Z. Chen, S. Tep, B. Ason, E. P. Somers, A. B. Sachs, S. R. Bartz, J. Tian, J. Chin, B. K. Hubbard, K. K. Wong, L. J. Mitaun, siRNA-induced liver ApoB knockdown lowers serum LDL-cholesterol in a mouse model with human-like serum lipids. *J. Lipid Res.* **52**, 1084–1097 (2011).
27. M. S. Hamza, C. Kumar, S. M. Chia, V. Anandalakshmi, N. Boo, W. Strapps, M. Robinson, M. Gaguyong, S. Bartz, M. Tadin-Strapps, A. van Gool, S.-J. Shih, Alterations in the hepatic transcriptional landscape after RNAi mediated ApoB silencing in cynomolgus monkeys. *Atherosclerosis* **242**, 383–395 (2015).
28. R. G. Lee, J. Crosby, B. F. Baker, M. J. Graham, R. M. Crooke, Antisense technology: An emerging platform for cardiovascular disease therapeutics. *J. Cardiovasc. Transl. Res.* **6**, 969–980 (2013).
29. J. J. P. Kastelein, M. K. Wedel, B. F. Baker, J. Su, J. D. Bradley, R. Z. Yu, E. Chuang, M. J. Graham, R. M. Crooke, Potent reduction of apolipoprotein B and low-density lipoprotein cholesterol by short-term administration of an antisense inhibitor of apolipoprotein B. *Circulation* **114**, 1729–1735 (2006).
30. E. A. Fisher, M. Pan, X. Chen, X. Wu, H. Wang, H. Jamil, J. D. Sparks, K. J. Williams, The triple threat to nascent apolipoprotein B. Evidence for multiple, distinct degradative pathways. *J. Biol. Chem.* **276**, 27855–27863 (2001).
31. H. N. Ginsberg, N. A. Le, J. C. Gibson, Regulation of the production and catabolism of plasma low density lipoproteins in hypertriglyceridemic subjects. Effect of weight loss. *J. Clin. Invest.* **75**, 614–623 (1985).
32. Y. Arad, R. Ramakrishnan, H. N. Ginsberg, Lovastatin therapy reduces low density lipoprotein apoB levels in subjects with combined hyperlipidemia by reducing the production of apoB-containing lipoproteins: Implications for the pathophysiology of apoB production. *J. Lipid Res.* **31**, 567–582 (1990).
33. J. L. Dixon, H. N. Ginsberg, Regulation of hepatic secretion of apolipoprotein B-containing lipoproteins: Information obtained from cultured liver cells. *J. Lipid Res.* **34**, 167–179 (1993).
34. A. Grefhorst, B. M. Elzinga, P. J. Voshol, T. Kok, V. W. Bloks, F. H. van der Sluijs, L. M. Havekes, J. A. Romijn, H. J. Verkade, F. Kuipers, Stimulation of lipogenesis by pharmacological activation of the liver X receptor leads to production of large, triglyceride-rich very low density lipoprotein particles. *J. Biol. Chem.* **277**, 34182–34190 (2002).
35. J. D. Brunzell, W. R. Hazzard, D. Porte Jr., E. L. Bierman, Evidence for a common, saturable, triglyceride removal mechanism for chylomicrons and very low density lipoprotein in man. *J. Clin. Invest.* **52**, 1578–1585 (1973).
36. D. W. Bilheimer, J. L. Goldstein, S. M. Grundy, Reduction in cholesterol and low density lipoprotein synthesis after portacaval shunt surgery in a patient with homozygous familial hypercholesterolemia. *J. Clin. Invest.* **56**, 1420–1430 (1975).
37. J. S. Millar, C. Maugeais, K. Ikewaki, D. M. Kolansky, P. H. R. Barrett, E. C. Budreck, R. C. Boston, N. Tada, S. Mochizuki, J. C. Defesche, J. M. Wilson, D. J. Rader, Complete deficiency of the low-density lipoprotein receptor is associated with increased apolipoprotein B-100 production. *Arterioscler. Thromb. Vasc. Biol.* **25**, 560–565 (2005).
38. K. Nagashima, C. Lopez, D. Donovan, C. Ngai, N. Fontanez, A. Bensadoun, J. Fruchart-Najib, S. Holleran, J. S. Cohn, R. Ramakrishnan, H. N. Ginsberg, Effects of the PPAR γ agonist pioglitazone on lipoprotein metabolism in patients with type 2 diabetes mellitus. *J. Clin. Invest.* **115**, 1323–1332 (2005).
39. J. S. Millar, G. Reyes-Soffer, P. Jumes, R. L. Dunbar, E. M. deGoma, A. L. Baer, W. Karmally, D. S. Donovan, H. Rafeek, L. Pollan, J. Tohyama, A. O. Johnson-Levonas, J. A. Wagner, S. Holleran, J. Obunike, Y. Liu, R. Ramakrishnan, M. E. Lassman, L. E. Gutstein, H. N. Ginsberg, D. J. Rader, Anacetrapib lowers LDL by increasing ApoB clearance in mildly hypercholesterolemic subjects. *J. Clin. Invest.* **125**, 2510–2522 (2015).
40. J. L. Dixon, S. Furukawa, H. N. Ginsberg, Oleate stimulates secretion of apolipoprotein B-containing lipoproteins from Hep G2 cells by inhibiting early intracellular degradation of apolipoprotein B. *J. Biol. Chem.* **266**, 5080–5086 (1991).
41. T. Ota, C. Gayet, H. N. Ginsberg, Inhibition of apolipoprotein B100 secretion by lipid-induced hepatic endoplasmic reticulum stress in rodents. *J. Clin. Invest.* **118**, 316–332 (2008).
42. P. Siri, N. Candela, Y.-L. Zhang, C. Ko, S. Eusufzai, H. N. Ginsberg, L.-S. Huang, Post-transcriptional stimulation of the assembly and secretion of triglyceride-rich apolipoprotein B lipoproteins in a mouse with selective deficiency of brown adipose tissue, obesity, and insulin resistance. *J. Biol. Chem.* **276**, 46064–46072 (2001).
43. F. Magkos, E. Fabbri, K. Korenblat, A. L. Okunade, B. W. Patterson, S. Klein, Reproducibility of glucose, fatty acid and VLDL kinetics and multi-organ insulin sensitivity in obese subjects with non-alcoholic fatty liver disease. *Int. J. Obes.* **35**, 1233–1240 (2011).
44. M. K. Hellerstein, R. A. Neese, Mass isotopomer distribution analysis at eight years: Theoretical, analytic, and experimental considerations. *Am. J. Physiol.* **276**, E1146–E1170 (1999).
45. R. Ramakrishnan, J. D. Ramakrishnan, Using mass measurements in tracer studies—a systematic approach to efficient modeling. *Metabolism* **57**, 1078–1087 (2008).
46. N. R. Matthán, M. Raeini-Sarjaz, A. H. Lichtenstein, L. M. Ausman, P. J. H. Jones, Deuterium uptake and plasma cholesterol precursor levels correspond as methods for measurement of endogenous cholesterol synthesis in hypercholesterolemic women. *Lipids* **35**, 1037–1044 (2000).
47. E. P. Lutz, M. Merkel, Y. Kako, K. Melford, H. Radner, J. L. Breslow, A. Bensadoun, I. J. Goldberg, Heparin-binding defective lipoprotein lipase is unstable and causes abnormalities in lipid delivery to tissues. *J. Clin. Invest.* **107**, 1183–1192 (2001).
48. J.-F. Hocquette, B. Graulet, T. Olivecrona, Lipoprotein lipase activity and mRNA levels in bovine tissues. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **121**, 201–212 (1998).
49. K.-I. Hirano, S. G. Young, R. V. Farese Jr., J. Ng, E. Sande, C. Warburton, L. M. Powell-Braxton, N. O. Davidson, Targeted disruption of the mouse *apobec-1* gene abolishes apolipoprotein B mRNA editing and eliminates apolipoprotein B48. *J. Biol. Chem.* **271**, 9887–9890 (1996).

Acknowledgments: We acknowledge laboratory assistance from M. Pavlyha and K. Poulos, The Research Pharmacy at Columbia University Medical Center, and the inpatient and outpatient nursing staff and Bionutrition unit of the ICTR at Columbia University Medical Center. We are grateful to N.R. Matthán and A.H. Lichtenstein (Tufts University) for providing the plasma sterol data. M. Graham (Ionis Pharmaceuticals) provided the apoB and MTP ASOs. N. Davidson (Washington University School of Medicine) provided the *Apobec-1* knockout mice. I. Goldberg (New York University School of Medicine) provided advice for the measurement of HL and LpL. R. M. Krauss (Children's Hospital Oakland Research Institute) provided assistance with the analysis of lipoprotein size/particle number and data interpretation. **Funding:** This study was funded by Sanofi-Genzyme. Additional support for this work was obtained from the National Center for Advancing Translational Sciences, NIH (through UL1 TR000040), and from the National Heart, Lung, and Blood Institute (through R01 HL55638). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. **Author contributions:** G.R.-S. designed and conducted all the human studies and participated in all aspects of writing the manuscript; B.M. designed and conducted all the cell experiments and wrote part of the manuscript; A.H.-O. designed and conducted all the mouse experiments and wrote part of the manuscript; M.D.-D. conducted the human studies; J.J. conducted the human studies; J.O. designed and conducted some of the human experiments; T.T. designed and conducted some of the human studies and contributed to the manuscript; C.N. conducted the human studies; N.F. conducted the human experiments; D.S.D. conducted the human studies; W.K. designed and conducted some of the human studies and wrote parts of the manuscript; S.H. designed and conducted the human studies; R.R. designed and conducted

RESEARCH ARTICLE

the human studies and wrote part of the manuscript; R.S.M. designed the human studies and wrote part of the manuscript; H.N.G. designed the human, cell, and mouse studies, conducted the human studies, and participated in all aspects of writing the manuscript. **Competing interests:** R.S.M. was an employee of Sanofi-Genzyme during the active study phase. G.R.-S. reports grants and nonfinancial support from Merck during the conduct of the study. D.S.D. reports grants from Merck during the conduct of the study. H.N.G. reports grants from Merck during the conduct of the study and personal consulting fees from Merck outside the submitted work. H.N.G. was also on the Sanofi advisory board during the time that this work was done. All other authors declare that they have no competing interests. **Data and materials availability:** Data are stored at Columbia and Sanofi-Genzyme. Data requests should be addressed to C. Sensinger of Genzyme (Charlotte.Sensinger@genzyme.com) or the study authors. Ionis Pharmaceuticals holds the patent on human apoB ASO (mipomersen), but mipomersen is now FDA-approved for the treatment of homozygous FH. Mouse apoB ASO was provided by Ionis Pharmaceuticals via a

material transfer agreement (MTA) with Columbia University. The human siRNA for *APOB* was provided by Merck Pharmaceutical via an MTA with Columbia University. As of December 2, 2015, Isis Pharmaceuticals Inc. has changed its name to Ionis Pharmaceuticals Inc.

Submitted 11 August 2015

Accepted 7 January 2016

Published 27 January 2016

10.1126/scitranslmed.aad2195

Citation: G. Reyes-Soffer, B. Moon, A. Hernandez-Ono, M. Dionizovik-Dimanovski, J. Jimenez, J. Obunike, T. Thomas, C. Ngai, N. Fontanez, D. S. Donovan, W. Karmally, S. Holleran, R. Ramakrishnan, R. S. Mittleman, H. N. Ginsberg, Complex effects of inhibiting hepatic apolipoprotein B100 synthesis in humans. *Sci. Transl. Med.* **8**, 323ra12 (2016).

Complex effects of inhibiting hepatic apolipoprotein B100 synthesis in humans

Gisette Reyes-Soffer, Byoung Moon, Antonio Hernandez-Ono, Marija Dionizovik-Dimanovski, Jhonsua Jimenez, Joseph Obunike, Tiffany Thomas, Colleen Ngai, Nelson Fontanez, Daniel S. Donovan, Wahida Karmally, Stephen Holleran, Rajasekhar Ramakrishnan, Robert S. Mittleman and Henry N. Ginsberg (January 27, 2016)

Science Translational Medicine **8** (323), 323ra12. [doi: 10.1126/scitranslmed.aad2195]

Editor's Summary

Making sense of antisense

Mipomersen is an FDA-approved antisense oligonucleotide that lowers low density lipoprotein (LDL) in patients with high cholesterol by targeting apolipoprotein B (apoB) synthesis. Although safe, how mipomersen works exactly in humans is unclear. Reyes-Soffer and colleagues found in healthy volunteers that the drug reduced levels of LDL and its precursor, very low density lipoproteins (VLDL), by increasing clearance of both of these vessel-clogging lipoproteins rather than reducing their secretion from the liver. The direct clearance of VLDL led to reduced production of LDL. Studies in mice and cell lines demonstrated how the liver compensates for reduced apoB synthesis to potentially avoid hepatic steatosis.

The following resources related to this article are available online at <http://stm.sciencemag.org>.
This information is current as of June 17, 2016.

Article Tools	Visit the online version of this article to access the personalization and article tools: http://stm.sciencemag.org/content/8/323/323ra12
Supplemental Materials	"Supplementary Materials" http://stm.sciencemag.org/content/suppl/2016/01/25/8.323.323ra12.DC1
Related Content	The editors suggest related resources on <i>Science</i> 's sites: http://stm.sciencemag.org/content/scitransmed/8/323/323rv2.full
Permissions	Obtain information about reproducing this article: http://www.sciencemag.org/about/permissions.dtl

Science Translational Medicine (print ISSN 1946-6234; online ISSN 1946-6242) is published weekly, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue, NW, Washington, DC 20005. Copyright 2016 by the American Association for the Advancement of Science; all rights reserved. The title *Science Translational Medicine* is a registered trademark of AAAS.