

# Arsenic in tube well water in Bangladesh: health and economic impacts and implications for arsenic mitigation

Sara V Flanagan,<sup>a</sup> Richard B Johnston<sup>b</sup> & Yan Zheng<sup>a</sup>

**Abstract** A national drinking water quality survey conducted in 2009 furnished data that were used to make an updated estimate of chronic arsenic exposure in Bangladesh. About 20 million and 45 million people were found to be exposed to concentrations above the national standard of 50 µg/L and the World Health Organization's guideline value of 10 µg/L, respectively. From the updated exposure data and all-cause mortality hazard ratios based on local epidemiological studies, it was estimated that arsenic exposures to concentrations > 50 µg/L and 10–50 µg/L account for an annual 24 000 and perhaps as many as 19 000 adult deaths in the country, respectively. Exposure varies widely in the 64 districts; among adults, arsenic-related deaths account for 0–15% of all deaths. An arsenic-related mortality rate of 1 in every 18 adult deaths could represent an economic burden of 13 billion United States dollars (US\$) in lost productivity alone over the next 20 years. Arsenic mitigation should follow a two-tiered approach: (i) prioritizing provision of safe water to an estimated 5 million people exposed to > 200 µg/L arsenic, and (ii) building local arsenic testing capacity. The effectiveness of such an approach was demonstrated during the United Nations Children's Fund 2006–2011 country programme, which provided safe water to arsenic-contaminated areas at a cost of US\$ 11 per capita. National scale-up of such an approach would cost a few hundred million US dollars but would improve the health and productivity of the population, especially in future generations.

Abstracts in **عربي**, **中文**, **Français**, **Русский** and **Español** at the end of each article.

## Introduction

Exposure to arsenic through drinking water sourced from groundwater is a global public health problem that is particularly devastating in Bangladesh.<sup>1,2</sup> According to survey data from 2000 to 2010, an estimated 35 to 77 million people in the country have been chronically exposed to arsenic in their drinking water in what has been described as the largest mass poisoning in history.<sup>2,3</sup> In rural areas, 97% of the population relies on tube wells<sup>4</sup> installed since the 1970s to reduce disease from ingestion of pathogen-laden surface waters. Unfortunately, this has resulted in a population highly exposed to arsenic but with limited means or incentives for seeking safe water alternatives. First detected in well water in the early 1990s, arsenic is released from sediment by biogeochemical processes that promote reducing environments.<sup>5,6</sup> The tube wells, affordably priced at about 100 United States dollars (US\$), draw the arsenic-containing groundwater from a shallow depth of 10–70 m.<sup>3</sup> Groundwater from depths > 150 m usually contains less arsenic<sup>3</sup> and can be a sustainable drinking water source.<sup>7</sup>

The health implications of chronic arsenic exposure in such a large population are substantial.<sup>2</sup> Between 2000 and 2003, 4.94 million tube wells throughout Bangladesh were tested for arsenic and marked as safe or unsafe.<sup>8,9</sup> Since then, well switching has partially succeeded in reducing exposure.<sup>10</sup> However, sustaining the behaviour change required for long-term sharing of wells is difficult. Additionally, severely affected areas have few if any safe water options and need alternative drinking water sources. Areas showing high proportions of unsafe wells (i.e. wells whose water contains arsenic in concentrations > 50 µg/L, the Bangladeshi drinking water standard) are largely the same areas experiencing the highest arsenic concentrations (often > 200 µg/L). This suggests that

interventions targeting areas with the highest proportion of unsafe wells are also likely to reach the population exposed to the highest arsenic concentrations and hence at highest risk of experiencing adverse health outcomes.<sup>11</sup> Mitigating the problem of water containing high levels of arsenic requires a sizeable investment in the water supply infrastructure. This paper provides evidence that such investment is economically justified when the health and economic burdens of unabated arsenic exposure are considered.

## Arsenic exposure from drinking water in 2009

The 2009 Bangladesh Multiple Indicator Cluster Survey (MICS) included collection of drinking water for arsenic tests from 15 000 randomized households nationwide.<sup>11</sup> The National Drinking Water Quality Survey report used an estimated national population of 164 million to estimate that 22 million and 5.6 million people are drinking water with arsenic concentrations > 50 µg/L and > 200 µg/L, respectively. According to preliminary census figures for 2011, the population of Bangladesh is about 142.3 million. Based on this figure, the people drinking water having arsenic concentrations > 50 µg/L and > 200 µg/L are approximately 19 million and 5 million, respectively. These estimates may be revised upwards when the final 2011 census figure is released. The proportion of water samples with arsenic in excess of permissible limits was found to be lower in the MICS survey than in previous national well surveys, which suggests important progress in mitigation (Table 1), although differences in sample collection (e.g. use of household drinking water versus source water) could also explain the difference.<sup>11</sup>

<sup>a</sup> Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, United States of America.

<sup>b</sup> Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland.

Correspondence to Yan Zheng (e-mail: yzheng@ldeo.columbia.edu).

(Submitted: 13 December 2011 – Revised version received: 12 August 2012 – Accepted: 20 August 2012 – Published online: 14 September 2012)

Table 1. Arsenic concentration in drinking water and proportions exposed as determined by testing during national surveys, Bangladesh

Arsenic concentration (µg/L)	BGS/DPHE 2000 (n = 3 534)		MICS 2009 (n = 14 442)	
	Proportion (%)	Cumulative (%)	Proportion (%)	Cumulative (%)
0–10	57.9	57.9	68.0	68.0
10.1–50	17.1	75.1	18.7	86.6
50.1–100	8.9	84.0	7.2	93.8
100.1–150	4.2	88.2	1.4	95.2
150.1–200	2.9	91.1	1.4	96.6
200.1–250	2.1	93.2	1.1	97.8
250.1–300	1.8	94.9	0.4	98.2
300+	5.1	100	1.8	100

BGS, British Geological Survey; DPHE, Department of Public Health Engineering; MICS, Multiple Indicator Cluster Survey.

## Modelling arsenic-related mortality

Chronic arsenic exposure is linked to a range of dose-dependent conditions, including cancers of the skin, bladder, kidney and lung,<sup>12–14</sup> as well as skin lesions, arterial hypertension and cardiovascular disease, pulmonary disease, peripheral vascular disease, diabetes mellitus and neuropathy.<sup>2</sup> In Bangladesh, the risk of dying from ingestion of arsenic in drinking water has been shown to depend on the level of arsenic exposure.<sup>15,16</sup> Sohel et al. analysed survival data for 1991–2000 from a health and demographic surveillance system covering 115 903 people in Matlab (Table 2). After adjusting for potential confounders such as age, sex, education and asset score (as an indicator of household wealth), they found that arsenic exposure through drinking water accounts for considerable excess mortality among adults in rural Bangladesh.<sup>15</sup>

The Health Effects of Arsenic Longitudinal Study (HEALS), which followed a cohort of 11 746 people in

Araihazar subdistrict from October 2000 to February 2009,<sup>16</sup> also showed that arsenic exposure is associated with a higher risk of death (Table 3). Although both Argos et al. and Sohel et al. found this increased risk even at low exposure levels (10–50 µg/L), historical exposure to concentrations > 50 µg/L arsenic may have introduced bias.<sup>17</sup> To reduce the risk of bias, the population exposed to 0–10 µg/L was used as a reference group, but because of uncertainties in lifetime exposure history in both studies, the dose category may have been assigned incorrectly, especially at the lower dose. Sohel et al. attempted to construct an exposure history for each subject but was unable to do so for those already deceased. Additionally, Sohel et al. found the hazard ratio (HR) to be higher for all non-accidental deaths than for any of the three known arsenic-related causes of death – cancer, cardiovascular problems, infection – at an exposure level of 10–50 µg/L (Table 2),<sup>15</sup> which suggests that factors other than arsenic exposure could have influenced the findings. Although the HRs from these studies

are fraught with uncertainties that bear further investigation, we used them to estimate arsenic-related mortality in Bangladesh because they were the best data available.

To assess the impact of arsenic exposure on mortality in Bangladesh, we calculated the excess deaths from the estimated risk of death (hazard) among adults in each arsenic exposure category (Table 2 and Table 3). The MICS 2009 drinking water quality survey provided the population exposure estimates.<sup>11</sup> From the resulting population attributable fraction (PAF) we estimated the annual number of deaths for each district by using the area's adult population (based on the census and the age distribution from the Bangladesh Demographic and Health Survey 2007)<sup>18</sup> and an estimate of the crude death rate. Because Bangladesh has no active vital registry system, we used a crude death rate for adults (> 15 years old) of 8.5 deaths per 1000 population, a figure based on WHO mortality estimates<sup>19</sup> and consistent with ICDDR,B Health and Demographic Surveillance System observations in Matlab and with crude death rates in other countries of southern Asia.

Using Sohel et al.'s HR for non-accidental deaths, we modelled excess deaths for all districts and arrived at an annual total of nearly 43 000 deaths, representing about 5.6% of all deaths, as being attributable to chronic arsenic exposure at current exposure levels (Table 4). On the basis of Sohel's cause-specific mortality HRs, about 1 in 16 cancer deaths, 1 in 36 cardiovascular disease deaths and 1 in 19 deaths from infections are attributable to arsenic exposure. We used Sohel et al.'s HR for non-accidental deaths because Argos et al.'s HR for the 10–50 µg/L exposure level is implausible, since it is

Table 2. Hazard ratios (HRs) for death from arsenic exposure, by cause of death and average arsenic concentration in drinking water, in a cohort of 115 903 people,<sup>15</sup> Bangladesh

Average arsenic concentration (µg/L)	HR for cause of death (95% CI)			
	Nonaccidental	Cancer	Cardiovascular	Infection
< 10 <sup>a</sup>	1.00	1.00	1.00	1.00
10–49	1.16 (1.06–1.26)	1.10 (0.77–1.59)	1.03 (0.82–1.29)	1.09 (0.92–1.30)
50–149	1.26 (1.18–1.36)	1.44 (1.06–1.95)	1.16 (0.96–1.40)	1.30 (1.13–1.49)
150–299	1.36 (1.27–1.47)	1.75 (1.28–2.40)	1.23 (1.01–1.51)	1.51 (1.31–1.75)
≥ 300	1.35 (1.23–1.48)	1.56 (1.06–2.30)	1.37 (1.07–1.77)	1.59 (1.33–1.91)

CI, confidence interval.

<sup>a</sup> Reference category.

Table 3. **Hazard ratios (HRs) for death from arsenic exposure, by cause of death and baseline arsenic concentration in drinking water, in a cohort of 11 746 people,<sup>16</sup> Bangladesh**

Baseline arsenic concentration ( $\mu\text{g/L}$ )	Cause of death	
	All-cause HR (95% CI)	Chronic disease HR (95% CI)
< 10.1	1.00	1.00
10.1–50.0	1.34 (0.99–1.82)	1.33 (0.94–1.87)
50.1–150.0	1.09 (0.81–1.47)	1.22 (0.87–1.70)
150.1–864.0	1.68 (1.26–2.23)	1.68 (1.21–2.33)

CI, confidence interval.

not significant at the 0.05 significance level, is higher than for the 50–150  $\mu\text{g/L}$  exposure group, and predicts nearly twice as many excess deaths as Sohel et al.'s HR (Table 5). Interestingly, under either study the excess deaths among people exposed to arsenic concentrations of 10–50  $\mu\text{g/L}$  (below the national standard) represent from 45% to 62% of all arsenic-related deaths. However, a proportion of the population that is currently in the 10–50  $\mu\text{g/L}$  exposure group may have been exposed to higher arsenic concentrations in the past and have an increased risk of death reflective of previous rather than current exposure. In light of this, we used the total number of arsenic-attributable deaths – about 43 000 deaths per year – for our economic impact assessment, since it more accurately reflects total exposure, past and present.

## Economic implications

We estimated the economic losses resulting from the arsenic-related mortality burden by calculating lost productivity in terms of per capita gross domestic product (GDP). According to estimates by the International Monetary Fund, the per capita GDP for Bangladesh in 2009 was 1465 purchasing power parity dollars. If we assume steady economic growth and an average loss of 10 years of productivity per arsenic-attributable death, over the next 20 years arsenic-related mortality in Bangladesh (1 of every 18 deaths) could lead to a loss of US\$ 12.5 billion, provided arsenic exposure (> 10  $\mu\text{g/L}$ ) remains the same as in 2009. We made this estimate using Sohel et al.'s HRs and a discount rate of 5%.<sup>20</sup> Our assumption of an average loss of 10 years of productivity per arsenic-attributable death was based on lost productivity owing to deaths from types of cancer known to be arsenic-related

and may be a conservative assumption because medical care capacity in Bangladesh is limited. The average person dying of cancer in the United States of America loses 15.4 years of life and, for the four types of cancer linked to arsenic exposure (skin, bladder, kidney and lung), the average loss ranges from 11 to 18 years.<sup>2,21</sup> Although life expectancy in the United States is higher than in Bangladesh, the proportion of time people spend working is probably higher in Bangladesh, so any loss of life in Bangladesh would translate into a greater reduction in lifetime productivity. Because the loss in GDP attributable to deaths does not take into account health costs or other costs to society, it probably underestimates the full economic burden. This burden can be expected to grow as the country develops and life expectancy rises. The morbidity burden will also increase as diagnostic tests improve and better treatment methods prolong the lives of people with chronic arsenic-related disease, and the costs of medical care will increase in tandem.

## Consequences of delaying action

In Bangladesh, arsenic-related diseases and deaths will increase in the future because the latency period after exposure lasts several decades.<sup>2</sup> Studies on chronic arsenic exposure in utero and in early childhood suggest an increased risk of fetal loss, infant death, reduced birth weight and impaired cognitive function in children, as well as significantly higher risks of impaired lung function, renal cancer and death from lung cancer, lung disease and acute myocardial infarction later in life.<sup>22–26</sup> Since an entire generation has now grown up exposed to arsenic, some children will become “arsenic orphans” as their caretakers suc-

cumb to arsenic-related diseases. These children may also be exposed to arsenic themselves, which would perpetuate the cycle of arsenic-related disease.

It is illustrative to examine the impact of arsenic exposure on children not yet born, whose future health will be affected by the concentration of arsenic in the water they begin drinking *in utero*, as shown by several studies.<sup>22–27</sup> We contemplate three scenarios for population exposure to arsenic in concentrations > 50  $\mu\text{g/L}$ : in the first and worst, exposure is constant beginning in 2000; in the second and best, exposure has been eliminated by 2010; in the third and most realistic, exposure is reduced to 13% by 2010 (as found in MICS 2009) and completely eliminated by 2030. How will these exposure scenarios affect today's children in the future? The proportion of eventual deaths attributable to arsenic exposure above the national standard in each year's birth cohort ranges from 0% when exposure to drinking water containing arsenic in concentrations > 50  $\mu\text{g/L}$  has been eliminated by the respective year, to 5.8% if the exposure level remains the same as in 2000. Overall, only 1.1% of eventual deaths in the 2000–2030 cohorts would be attributable to arsenic if exposure to concentrations > 50  $\mu\text{g/L}$  had been eliminated by 2010. However, if exposure levels throughout 2000–2030 were to remain the same as in 2000, 5.8% of all eventual deaths in the 2000–2030 cohort would be attributable to arsenic. The most likely scenario will lie in between: if exposure to arsenic in concentrations > 50  $\mu\text{g/L}$  is eliminated by 2030, 2.4% of the cohort's future deaths will be attributable to arsenic. In absolute terms, if about 90 million children are born between 2000 and 2030, between 1 and 5 million of their eventual deaths will be attributable to exposure to arsenic concentrations above the national standard, depending on the exposure scenario. This exercise shows that any population-level reduction in arsenic exposure will result in decreased arsenic-related morbidity and mortality among children yet to be born. Similarly, any failure to sustain progress in arsenic mitigation will result in deaths that could have been prevented among members of future generations. However, because of uncertainty and individual variation in arsenic exposure and the latency period before disease onset, these analyses are qualitative and semiquantitative predictions at best.

Table 4. Population attributable fraction (PAF) of deaths from arsenic exposure and arsenic-attributable excess deaths (ED) per year for different arsenic concentrations in drinking water, by district, Bangladesh, 2011

District	2011 adult population (thousands)	PAF (Sohel et al.) <sup>15</sup>	PAF (Argos et al.) <sup>16</sup>	Annual no. of arsenic-attributable ED by arsenic concentration (in µg/L) and in total (Sohel et al.) <sup>15</sup>				ED (Argos et al.) <sup>16</sup>	
				10–49	50–149	150–299	≥ 300		Total deaths
Bagerhat	931	0.036	0.045	133	103	41	8	285	361
Bandarban	244	0.013	0.024	28	0	0	0	28	51
Barguna	562	0.003	0.005	13	0	0	0	13	23
Barisal	1459	0.019	0.031	158	24	20	40	241	392
Bhola	1120	0.003	0.006	32	0	0	0	32	59
Bogra	2147	0.022	0.034	293	90	29	0	412	619
Brahmanbaria	1789	0.090	0.116	240	373	273	485	1371	1765
Chandpur <sup>a</sup>	1524	0.115	0.152	77	324	461	640	1503	1976
Chittagong <sup>a</sup>	4783	0.041	0.062	1140	383	98	72	1694	2514
Chuadanga	715	0.087	0.116	296	173	49	12	531	708
Comilla <sup>a</sup>	3379	0.130	0.162	356	1066	1262	1064	3748	4673
Cox's Bazar	1449	0.004	0.006	34	13	0	0	47	68
Dhaka	7564	0.020	0.026	555	435	305	0	1295	1662
Dinajpur	1892	0.007	0.013	115	0	0	0	115	212
Faridpur	1189	0.133	0.171	511	439	292	112	1354	1737
Feni	905	0.088	0.084	177	392	88	21	678	649
Gaibandha	1496	0.071	0.107	633	189	49	32	902	1364
Gazipur	2123	0.006	0.009	63	23	30	0	116	170
Gopalganj	732	0.145	0.171	223	341	267	73	904	1069
Habiganj	1312	0.089	0.115	542	356	101	0	999	1294
Jamalpur	1443	0.015	0.063	76	0	0	0	76	781
Jessore	1747	0.040	0.120	380	86	32	0	498	1785
Jhalokathi	380	0.096	0.038	552	511	303	66	1431	123
Jhenaidah	1119	0.023	0.104	65	10	0	0	75	991
Joypurhat	579	0.077	0.028	449	247	43	0	739	140
Khagrachari	387	0.002	0.003	4	3	0	0	7	8
Khulna	1461	0.056	0.076	414	214	49	19	696	952
Kishoreganj	1817	0.085	0.107	659	489	123	44	1315	1666
Kurigram	1306	0.048	0.079	446	71	23	0	540	883
Kushtia	1231	0.050	0.076	407	119	0	0	526	796
Lakshmipur	1090	0.090	0.119	343	251	77	165	835	1106
Lalmonirhat	796	0.017	0.031	113	0	0	0	113	209
Madaripur	732	0.119	0.154	246	223	212	61	742	961
Magura	582	0.083	0.104	204	157	20	30	410	515
Manikganj	878	0.093	0.116	363	275	30	30	698	870
Maulvibazar	1212	0.097	0.096	208	76	44	17	345	989
Meherpur	415	0.080	0.143	403	360	68	0	830	507
Munshiganj	905	0.066	0.073	85	221	120	85	511	561
Mymensingh	3212	0.047	0.074	1018	239	22	21	1301	2035
Naogaon	1641	0.024	0.043	322	10	0	0	332	597
Narail	455	0.092	0.111	116	136	97	10	360	432
Narayanganj	1845	0.047	0.055	160	269	306	0	735	870
Narsingdi	1403	0.052	0.066	236	217	87	85	626	788
Natore	1080	0.004	0.006	28	11	0	0	39	57
Nawabganj	1041	0.045	0.055	169	161	74	0	403	487
Netrokona	1406	0.115	0.136	570	585	188	43	1386	1636
Nilphamari	1159	0.029	0.053	288	0	0	0	288	530
Noakhali <sup>a</sup>	1957	0.117	0.138	415	740	441	365	1962	2305
Pabna	1591	0.042	0.059	295	144	127	0	566	795
Panchagarh	625	0.017	0.032	93	0	0	0	93	170

(continues. . .)

(. . .continued)

District	2011 adult population (thousands)	PAF (Sohel et al.) <sup>15</sup>	PAF (Argos et al.) <sup>16</sup>	Annual no. of arsenic-attributable ED by arsenic concentration (in µg/L) and in total (Sohel et al.) <sup>15</sup>				ED (Argos et al.) <sup>16</sup>	
				10–49	50–149	150–299	≥ 300		Total deaths
Patuakhali	966	0.023	0.042	188	0	0	0	188	346
Pirojpur	703	0.044	0.074	232	23	8	0	263	447
Rajbari	662	0.021	0.070	174	99	0	23	295	397
Rajshahi	1639	0.051	0.028	150	81	46	11	288	395
Rangamati	380	0.014	0.025	44	0	0	0	44	81
Rangpur	1826	0.023	0.040	333	30	0	0	364	625
Satkhira	1257	0.054	0.149	196	107	0	36	339	1604
Shariatpur	730	0.110	0.074	465	317	311	93	1186	459
Sherpur	850	0.055	0.063	718	184	18	0	920	455
Sirajganj	1957	0.039	0.085	219	37	24	0	281	1423
Sunamganj <sup>a</sup>	1556	0.151	0.146	659	1192	127	34	2012	1936
Sylhet	2168	0.054	0.057	370	513	90	25	999	1063
Tangail	2275	0.040	0.064	621	118	30	0	769	1236
Thakurgaon	879	0.003	0.006	26	0	0	0	26	48
<b>Total</b>	<b>90657</b>	<b>0.056</b>	<b>0.074</b>	<b>19140</b>	<b>13250</b>	<b>6504</b>	<b>3823</b>	<b>42717</b>	<b>56425</b>

<sup>a</sup> One of five districts having the highest number of arsenic-attributable deaths per year (based on Soheli et al.'s hazard ratios).

## Mitigation strategy

According to the model, Comilla is the district with the highest number of arsenic-related deaths – 3748 adult deaths in 2009. This is because many people there are exposed to high arsenic concentrations (Table 4). Resulting losses in productivity could amount to US\$ 1.1 billion over the next 20 years in Comilla alone.<sup>20</sup> Supplying safe water to the district's population by installing water points with no more than 50 people per water point, as well as small communal piped water systems serving a few hundred households, would cost approximately US\$ 44.2–49.2 million depending on the choice of water supply technology.<sup>20</sup> This would be a fraction of the economic losses that would result from continued arsenic exposure, and the health benefits to generations not yet born would be incalculable. Despite the considerable capital costs involved, the benefits of an immediate investment in an improved water supply system would far outweigh the costs. Sustainability and appropriateness for a given setting should drive the choice of one arsenic mitigation technology over another.<sup>20</sup>

The water sector in Bangladesh urgently needs to find a sustainable way to supply safe water to people in areas with high arsenic exposure and to build capacity for local arsenic testing for surveillance.<sup>28</sup> Because of the dose–response relationship that characterizes arsenic-related health problems, the public health benefits of new safe water

supplies can be maximized by targeting grossly contaminated areas (i.e. with concentrations > 200 µg/L) first. Such areas are usually the ones having the highest proportion of wells with water that has arsenic concentrations > 50 µg/L. The Department of Public Health Engineering (DPHE) of Bangladesh and the United Nations Children's Fund (UNICEF) have succeeded in increasing access to safe water in Comilla for a per capita cost of only US\$ 11 by following this approach. Complete coverage of Comilla with safe water could be achieved for an additional US\$ 32 million. Thanks to the provision of safe water points in communities at risk as well as public education and social mobilization, the population drinking arsenic-safe water in the intervention area in Comilla increased steadily from 75% in 2007 to 81% in 2009.<sup>29</sup> However, in control areas access to arsenic-safe water decreased from 93% in 2007 to 83% in 2009, perhaps because of the continued installation of new and inexpensive but contaminated shallow tube wells and because adherence to well switching has declined as memories of arsenic awareness-raising activities have begun to fade. The greatest improvements in access were achieved among the poorest population quintiles in intervention areas, which points to success in targeting people living in poverty and extreme poverty.

As these examples suggest, past achievements can be lost if arsenic mitigation efforts are not sustained. Markings on wells from previous testing campaigns

have now worn off and the motivation for promoting arsenic-safe water has waned. The top-down blanket testing approach of the past left no infrastructure in place for monitoring existing wells or for testing new wells.<sup>30</sup> Building testing capacity locally will lead to sustained awareness in areas with high arsenic exposure and give people more control over their water supply, although instilling a social norm of periodically testing well water is essential for sustainability. Implementing a local pay-for-use testing system has already been found effective at motivating households to test wells and, in turn, has strengthened the commitment of the local population to undertake arsenic mitigation measures. By making it possible for people to know which local wells are contaminated and which ones are safe<sup>31</sup> and by strategically providing new water supply systems to the populations most exposed to arsenic, compliance with the national drinking water arsenic standard can be facilitated. Progress will not be even, however, since some areas will prove more challenging than others.<sup>32</sup> Social acceptability and sustainability are crucial factors to be considered when choosing among arsenic mitigation strategies, in addition to the costs of the technologies involved.<sup>20</sup> For example, technologies for removing arsenic from contaminated water would cost an average of four times as much over a span of 20 years as delivery of safe water obtainable from other sources, and would require high maintenance. Thus, technologies that avoid arsenic contami-



Table 5. **Population attributable fraction (PAF) of deaths and excess deaths (ED) from arsenic exposure based on hazard ratios from two published sources, Bangladesh**

Arsenic concentration ( $\mu\text{g/L}$ )	Percentage exposed	PAF (Sohel et al.) <sup>15</sup>	PAF (Argos et al.) <sup>16</sup>	ED (Sohel et al.) <sup>15</sup>	ED (Argos et al.) <sup>16</sup>
< 10	68.0	0	0	0	0
10.1–50	18.7	0.138	0.254	19 140	35 210
50.1–150	8.6	0.206	0.083	13 250	5302
150.1–300	3.0	0.265	0.405	6504	9945
> 300	1.8	0.259	0.405	3823	5969
<b>Total</b>	<b>100.0</b>	<b>0.056</b>	<b>0.074</b>	<b>42 717</b>	<b>56 425</b>
> 50 $\mu\text{g/L}$ only	13.4	0.030	0.027	23 577	21 215

nation, rather than remove arsenic, are more cost-effective in the long term.<sup>33</sup>

## Conclusion

In Bangladesh, ongoing exposure to arsenic in drinking water calls for renewed and sustained mitigation efforts. Exposure to arsenic could be eliminated by 2030 if the government invested a small fraction of its annual GDP growth in providing an arsenic-safe water supply and improving water quality monitoring and surveillance activities. Reductions in arsenic-related mortality would be noted

within about 40 years, as suggested by observations in similarly exposed populations in Chile and, Taiwan (China), where arsenic-related cancer mortality started to decline gradually about 20 or 25 years after measures to reduce exposure were initiated and coronary heart disease mortality declined even faster.<sup>24,34–37</sup> The current generation may face the latent effects of lifetime exposure to arsenic even after switching to a safe water source, but for future generations, arsenic-attributable disease and death would be a thing of the past. If, on the other hand, population-wide chronic

arsenic exposure is allowed to continue unchecked or to worsen as the population grows and installs more private tube wells, future generations will be saddled with enormous health and productivity costs. Appropriate interventions and robust investments, if undertaken now, can prevent this from happening. ■

## Acknowledgements

This paper reflects the views of the authors only and not those of UNICEF. The authors thank Astrid van Agthoven, Syed Adnan Ibna Hakim, Mi Hua, Deqa Ibrahim Musa, Hans Spruijt and Siping Wang of UNICEF Bangladesh, as well as the Bangladesh Bureau of Statistics and the Department of Public Health and Engineering. This work was conducted when all authors were affiliated with the Water, Environmental Sanitation Section of UNICEF Bangladesh. Yan Zheng is also currently affiliated with Queens College, City University of New York.

**Funding:** UKAid, UNICEF and the Government of Bangladesh funded the SHEWA-B project and the MICS Water Quality Survey.

**Competing interests:** None declared.

## ملخص

### الزرنخ في مياه الآبار الأنبوبية في بنغلاديش: الآثار والتداعيات الصحية والاقتصادية المترتبة على التخفيف من

#### مضار الزرنخ

ويمكن أن يمثل معدل الوفيات المرتبط بالزرنخ المقدّر بحالة بين كل 18 حالة وفاة من البالغين عبئاً اقتصادياً قدره 13 مليار دولار أمريكي في الإنتاجية المفقودة وحدها على مدار العشرين سنة القادمة. ويجب أن يتبع التخفيف من مضار الزرنخ نهجاً من مستويين: (1) منح الأولوية لتوفير المياه النقية إلى 5 مليون شخص معرضين إلى تركيز أعلى من 200 ميكروجرام/لتر من الزرنخ، و(2) بناء قدرات اختبار محلية للزرنخ. وقد اتضحت فعالية هذا النهج أثناء البرنامج القطري لليونيسيف في الفترة من 2006 إلى 2011 الذي وفر المياه النقية للمناطق الملوثة بالزرنخ بتكلفة 11 دولاراً أمريكياً للفرد. وسيتكلف تحسين هذا النهج على الصعيد الوطني بضعة مئات الملايين من الدولارات الأمريكية غير أنه سيحسن صحة السكان وإنتاجيتهم، ولا سيما في الأجيال المستقبلية.

قدم استبيان وطني لنوعية مياه الشرب أجري في عام 2009 البيانات التي استخدمت لإجراء تقدير محدث للتعرض المزمن للزرنخ في بنغلاديش. وقد تبين تعرض حوالي 20 مليون و45 مليون شخص إلى تركيزات أعلى من المعيار الوطني 50 ميكروجرام/لتر ومن القيمة الإرشادية لمنظمة الصحة العالمية 10 ميكروجرام/لتر، على التوالي. وتم التوصل من بيانات التعرض المحدثة ونسب مخاطر الوفاة الناتجة عن جميع الأسباب المستندة على الدراسات الوبائية المحلية إلى تقدير يفيد بأن التعرض للزرنخ بتركيزات أعلى من 50 ميكروجرام/لتر و10–50 ميكروجرام/لتر يتسبب في حدوث 24000 حالة وفاة وعدد محتمل يصل إلى 19000 حالة وفاة بين البالغين سنوياً في البلد، على التوالي. ويختلف التعرض على نطاق واسع في الأربع والسنتين منطقتي؛ وتبلغ نسبة الوفيات المرتبطة بالزرنخ بين البالغين ما بين 0٪ إلى 15٪ من جميع حالات الوفاة.

## 摘要

### 孟加拉国民用井水中的砷：控制砷暴露对健康和经济的影响和意义

使用2009年孟加拉国全国饮用水水质调查的数据，对孟加拉国慢性砷暴露状况作出最新评估。据发现，分别有2000万和4500万人口饮用水砷浓度超过50  $\mu\text{g/L}$ 的

国家标准和10  $\mu\text{g/L}$ 的世界卫生组织指导值。基于这些最新砷暴露数据，结合当地人群由饮用水砷暴露导致的死亡风险率的流行病学研究成果，可估算到砷暴露浓度>

50  $\mu\text{g/L}$ 和10~50  $\mu\text{g/L}$ 分别造成每年2.4 万以及可能多达1.9 万的成人死亡。64 个省份的差异很大；在成年人中，砷相关的死亡占所有死亡人数的0%至15%。由于每18 个死亡的成年人中就有1 个与砷相关的死亡使生产力下降，这造成相当于未来20 年130 亿美元 (US\$) 国民生产总值的经济损失。降低砷暴露可两条路线并进：(一) 优先为饮用水砷浓度 > 200  $\mu\text{g/L}$ 的约500 万暴露人口提

供安全的饮用水，及(二) 建立地方砷检测能力。此方法的有效性在联合国儿童基金 2006-2011 年孟加拉国计划中得以示范，以人均11 美元的成本为砷危害地区安全供水。在孟加拉国全国范围推广这样的方案将花费数亿美元，但会提高国民的健康状况和避免国民生产力的损失，造福子孙后代。

## Résumé

### Arsenic dans l'eau des puits tubulaires au Bangladesh: impacts sanitaires et économiques, et implications en vue de sa réduction

Une enquête nationale sur la qualité de l'eau potable, menée en 2009, a fourni des données qui ont été utilisées pour faire une estimation actualisée de l'exposition chronique à l'arsenic au Bangladesh. On a découvert qu'environ 20 et 45 millions de personnes étaient exposées à des concentrations supérieures à la norme nationale de 50  $\mu\text{g/l}$  et à la valeur indicative de 10  $\mu\text{g/l}$  de l'Organisation mondiale de la Santé, respectivement. À partir des données d'exposition actualisées et des ratios de risque de mortalité toutes causes confondues, sur la base d'études épidémiologiques locales, on a estimé que l'exposition à des concentrations d'arsenic supérieures à 50  $\mu\text{g/l}$  et entre 10-50  $\mu\text{g/l}$  étaient à l'origine de 24 000, voire jusqu'à 19 000 décès annuels d'adultes dans le pays, respectivement. L'exposition varie considérablement dans les 64 districts. Chez les adultes, les décès associés à l'arsenic représentaient 0% à 15% de tous les décès. Un taux de mortalité associé

à l'arsenic de 1 pour 18 décès d'adultes pourrait représenter au cours des 20 prochaines années un fardeau économique de 13 milliards de dollars américains (USD), en termes de perte de productivité seulement. La diminution de la quantité d'arsenic doit suivre une approche à deux niveaux: (i) considérer comme prioritaire la fourniture d'eau potable à environ 5 millions de personnes exposées à plus de 200  $\mu\text{g/l}$  d'arsenic, et (ii) générer la capacité locale à effectuer des tests sur la teneur en arsenic. L'efficacité d'une telle approche a été démontrée lors du programme national 2006-2011 du Fonds des Nations unies pour l'enfance, qui a fourni de l'eau potable à des zones contaminées par l'arsenic, pour un coût de 11 USD par habitant. L'application à l'échelle nationale d'une telle approche coûterait quelques centaines de millions de dollars, mais permettrait d'améliorer la santé et la productivité de la population, en particulier pour les générations futures.

## Резюме

### Мышьяк в колодезной воде в Бангладеш: влияние на здоровье и экономику и предпосылки уменьшения вредного воздействия мышьяка

Для оценки длительного воздействия мышьяка в Бангладеш были использованы данные национального исследования качества питьевой воды, проведенный в 2009 г. Около 20 и 45 миллионов человек оказались подвержены воздействию концентраций мышьяка, превышающих национальный стандарт, равный 50  $\text{мкг/л}$ , и нормативное значение Всемирной организации здравоохранения, равное 10  $\text{мкг/л}$ , соответственно. Из уточненных данных воздействия и соотношений риска общей смертности, полученных из локальных эпидемиологических исследований, было установлено, что воздействие мышьяка в концентрациях > 50  $\text{мкг/л}$  и 10–50  $\text{мкг/л}$  является причиной соответственно 24 000 и, возможно, не менее 19 000 смертей среди взрослого населения в стране. Степень воздействия колеблется в широких пределах в 64 районах: среди взрослого населения число смертей, связанных с воздействием мышьяка, составляет 0–15% от общего числа смертей. Уровень смертности от мышьяка, составляющий 1 случай на каждые 18 смертей среди

взрослого населения, может стать причиной экономического ущерба в размере 13 млрд. долл. США только в результате снижения производительности труда в течении последующих 20 лет. Для уменьшения воздействия мышьяка следует придерживаться двухъярусного подхода: (i) отдавая приоритет снабжению безопасной водой примерно 5 млн. человек, подверженных концентрации мышьяка > 200  $\text{мкг/л}$ , и (ii) открывая местные учреждения для лабораторных исследований мышьяка. Эффективность подобного подхода была продемонстрирована при осуществлении Государственной программы Детского фонда ООН в 2006–2011 гг., в рамках которой зараженные мышьяком районы обеспечивались чистой водой, и расходы на которую составили 11 долл. США на душу населения. Расширение подобного подхода в масштабе страны стоило бы несколько млн. долл. США, но улучшило бы здоровье и производительность труда населения, особенно будущих поколений.

## Resumen

### El arsénico en el agua procedente de pozos entubados en Bangladesh: El impacto sobre la salud y la economía y las implicaciones en la mitigación de los efectos del arsénico

Una encuesta nacional sobre la calidad del agua potable llevada a cabo en 2009 proporcionó los datos que se emplearon para realizar un cálculo actualizado de la exposición crónica al arsénico en Bangladesh. Se descubrió que aproximadamente 20 millones de personas están expuestas a concentraciones superiores al nivel nacional de 50  $\mu\text{g/L}$  y unos 45 millones a concentraciones por encima del valor de referencia de la Organización Mundial de la Salud de 10  $\mu\text{g/L}$ . Partiendo de los datos

actualizados sobre la exposición y los cocientes de riesgo de mortalidad por todas las causas basados en estudios epidemiológicos, se calculó que la exposición al arsénico en concentraciones superiores a 50  $\mu\text{g/L}$  y entre 10–50  $\mu\text{g/L}$  causan, respectivamente, 24 000 y hasta 19 000 muertes entre los adultos en el país. La exposición varía mucho en los 64 distritos. Entre las personas adultas, los fallecimientos relacionados con el arsénico explican entre el 0 y el 15% de todas las muertes. La tasa

de mortalidad por arsénico de 1 de cada 18 adultos podría representar una carga económica de 13 mil millones de dólares estadounidenses (US\$) en productividad perdida en los próximos 20 años. La mitigación de los efectos del arsénico debería seguir un enfoque de dos niveles: (i) priorizar el suministro de agua salubre a aproximadamente 5 millones de personas expuestas a niveles de arsénico superiores a 200 µg/L, y (ii) construir centros locales con capacidad para realizar análisis de

arsénico. La eficacia de tal enfoque quedó demostrada durante el programa por países 2006–2011 del Fondo de las Naciones Unidas para la infancia, que suministró agua salubre a las áreas contaminadas con arsénico con un coste de US\$ 11 per cápita. La ampliación de semejante enfoque a nivel nacional costaría algunos cientos millones de dólares estadounidenses, pero mejoraría la salud y productividad de la población, sobre todo de las generaciones futuras.

## References

- Ravenscroft P, Brammer H, Richards K. *Arsenic pollution: a global synthesis*. Oxford: John-Wiley and Sons; 2009.
- Smith AH, Lingas E, Rahman M. Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. *Bull World Health Organ* 2000;78:1093–103. PMID:11019458
- Kinniburgh DG, Smedley PL, editors. *Arsenic contamination of groundwater in Bangladesh*. Keyworth: British Geological Survey; 2001.
- Multiple Indicator Cluster Survey (MICS) 2009. New York: United Nations Children's Fund, Bangladesh & Bangladesh Bureau of Statistics; 2010. Available from: [http://www.unicef.org/bangladesh/knowledgecentre\\_6292.htm](http://www.unicef.org/bangladesh/knowledgecentre_6292.htm) [accessed 21 August 2012]
- Nickson R, McArthur J, Burgess W, Ahmed KM, Ravenscroft P, Rahman M. Arsenic poisoning of Bangladesh groundwater. *Nature* 1998;395:338. doi:10.1038/26387 PMID:9759723
- Zheng Y, Stute M, van Geen A, Gavrieli I, Dhar R, Simpson HJ et al. Redox control of arsenic mobilization in Bangladesh groundwater. *Appl Geochem* 2004;19:201–14. doi:10.1016/j.apgeochem.2003.09.007
- Radloff KA, Zheng Y, Michael HA, Stute M, Bostick BC, Mihajlov I et al. Arsenic migration to deep groundwater in Bangladesh influenced by adsorption and water demand. *Nat Geosci* 2011;4:793–8. doi:10.1038/ngeo1283 PMID:22308168
- Towards an arsenic safe environment in Bangladesh: World Water Day, 22 March 2010*. Dhaka: United Nations, Bangladesh and the Government of Bangladesh; 2010. Available from [http://www.unicef.org/bangladesh/knowledgecentre\\_6131.htm](http://www.unicef.org/bangladesh/knowledgecentre_6131.htm) [accessed 21 August 2012].
- Johnston RB, Sarker M. Arsenic mitigation in Bangladesh: national screening data and case studies in three upazilas. *Journal of Environmental Science and Health* 2007;42:1889–96. PMID:17952790
- Van Geen A. Promotion of well-switching to mitigate the current arsenic crisis in Bangladesh. *Bull World Health Organ* 2002;80:732–7. PMID:12378292
- Bangladesh national drinking water quality survey 2009*. Dhaka: United Nations Children's Fund, Bangladesh; 2011. Available from: [http://www.unicef.org/bangladesh/knowledgecentre\\_6868.htm](http://www.unicef.org/bangladesh/knowledgecentre_6868.htm) [accessed 21 August 2012].
- Chen CJ, Kuo T, Wu M. Arsenic and cancers. *Lancet* 1988;331:414–5. doi:10.1016/S0140-6736(88)91207-X PMID:2893213
- Chen CJ, Chen C, Wu M, Kuo T. Cancer potential in liver, lung, bladder and kidney due to ingested inorganic arsenic in drinking water. *Br J Cancer* 1992;66:888–92. doi:10.1038/bjc.1992.380 PMID:1419632
- Smith AH, Goycolea M, Haque R, Biggs M. Marked increase in bladder and lung cancer mortality in a region of northern Chile due to arsenic in drinking water. *Am J Epidemiol* 1998;147:660–9. doi:10.1093/oxfordjournals.aje.a009507 PMID:9554605
- Sohel N, Persson L, Rahman M, Streatfield P, Yunus M, Ekstrom E-C et al. Arsenic in drinking water and adult mortality: a population-based cohort study in rural Bangladesh. *Epidemiology* 2009;20:824–30. doi:10.1097/EDE.0b013e3181bb56ec PMID:19797964
- Argos M, Kalra T, Rathouz P, Chen Y, Pierce B, Parvez F et al. Arsenic exposure from drinking water and all-cause and chronic-disease mortalities in Bangladesh (HEALS): a prospective cohort study. *Lancet* 2010;376:252–8. doi:10.1016/S0140-6736(10)60481-3 PMID:20646756
- Ahsan H, Chen Y, Parvez F, Zablotska L, Argos M, Hussain I et al. Arsenic exposure from drinking water and risk of premalignant skin lesions in Bangladesh: baseline results from the Health Effects of Arsenic Longitudinal Study. *Am J Epidemiol* 2006;163:1138–48. doi:10.1093/aje/kwj154 PMID:16624965
- Bangladesh Demographic and Health Survey 2007. Dhaka: National Institution for Population Research and Training; 2009.
- Mortality summary estimates for WHO Member States. In: *Global burden of disease estimates 2008*. Geneva: World Health Organization; 2011.
- Making economic sense for arsenic mitigation: a case study of Comilla district*. Dhaka: United Nations Children's Fund, Bangladesh; 2011. Available from: [http://www.unicef.org/bangladesh/knowledgecentre\\_6872.htm](http://www.unicef.org/bangladesh/knowledgecentre_6872.htm) [accessed 21 August 2012].
- SEER cancer statistics review 1975–2006*. Bethesda: National Cancer Institute; 2006. Available from: [http://seer.cancer.gov/csr/1975\\_2006/results\\_merged/topic\\_year\\_lost.pdf](http://seer.cancer.gov/csr/1975_2006/results_merged/topic_year_lost.pdf) [accessed 21 August 2012].
- Smith AH, Steinmaus C. Health effects of arsenic and chromium in drinking water: recent human findings. *Annu Rev Public Health* 2009;30:107–22. doi:10.1146/annurev.publhealth.031308.100143 PMID:19012537
- Dauphiné DC, Ferreccio C, Guntur S, Yuan Y, Hammond S, Balmes J et al. Lung function in adults following in utero and childhood exposure to arsenic in drinking water: preliminary findings. *Int Arch Occup Environ Health* 2011;84:591–600. doi:10.1007/s00420-010-0591-6 PMID:20972800
- Yuan Y, Marshall G, Ferreccio C, Steinmaus C, Liaw J, Bates M et al. Kidney cancer mortality: fifty-year latency patterns related to arsenic exposure. *Epidemiology* 2010;21:103–8. doi:10.1097/EDE.0b013e3181c21e46 PMID:20010213
- Smith AH, Marshall G, Yuan Y, Ferreccio C, Liaw J, von Ehrenstein O et al. Increased mortality from lung cancer and bronchiectasis in young adults after exposure to arsenic in utero and in early childhood. *Environ Health Perspect* 2006;114:1293–6. doi:10.1289/ehp.8832 PMID:16882542
- Yuan Y, Marshall G, Ferreccio C, Steinmaus C, Selvin S. Acute myocardial infarction mortality in comparison with lung and bladder cancer mortality in arsenic-exposed region II of Chile from 1950 to 2000. *Am J Epidemiol* 2007;166:1381–91. doi:10.1093/aje/kwm238 PMID:17875584
- Gardner R, Hamadani J, Grandér M, Tofail F, Nermell B, Palm B et al. Persistent exposure to arsenic via drinking water in rural Bangladesh despite major mitigation efforts. *Am J Public Health* 2011;101(Suppl 1):S333–8. doi:10.2105/AJPH.2010.300025 PMID:21778503
- George CM, Zheng Y, Graziano JH, Hossain Z, Rasul SB, Mey JL et al. Evaluation of an arsenic test kit for rapid well screening in Bangladesh. *Environ Sci Technol* 2012. Epub ahead of print 25 Sep 2012 doi:10.1021/es300253p PMID:22866936
- SHEWA-B health impact study midline report, November 2009*. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh; 2009.
- Zheng Y, Ravenscroft P, Rahman SM, Hakim SAI. Pay-for-use arsenic testing: promoting demand-driven mitigation and monitoring in Bangladesh. In: *Proceedings of the International Conference on Arsenic in the Environment*; July 22–27 2012; Cairns, Australia.
- Madajewicz M, Pfaff A, van Geen A, Graziano J, Hussein I, Momotaj H et al. Can information alone change behavior? Response to arsenic contamination of groundwater in Bangladesh. *J Dev Econ* 2007;84:731–54. doi:10.1016/j.jdeveco.2006.12.002
- Chen Y, van Geen AF, Graziano JH, Pfaff A, Madajewicz M, Parvez F et al. Reduction in urinary arsenic levels in response to arsenic mitigation efforts in Arai-hazar, Bangladesh. *Environ Health Perspect* 2007;115:917–23. doi:10.1289/ehp.9833 PMID:17589600
- Johnston RB, Hanchett S, Khan MH. The socio-economics of arsenic removal. *Nat Geosci* 2010;3:2–3. doi:10.1038/ngeo735
- Yang C-Y, Chiu H-F, Wu T-N, Chuang H-Y, Ho S-C. Reduction in kidney cancer mortality following installation of a tap water supply system in an arsenic-endemic area of Taiwan. *Arch Environ Health* 2004;59:484–8. doi:10.1080/00039890409603430 PMID:16381491
- Chiu H-F, Ho S-C, Yang C-Y. Lung cancer mortality reduction after installation of tap-water supply system in an arseniasis-endemic area in southwestern Taiwan. *Lung Cancer* 2004;46:265–70. doi:10.1016/j.lungcan.2004.05.012 PMID:15541810
- Marshall G, Ferreccio C, Yuan Y, Bates M, Steinmaus C, Selvin S et al. Fifty-year study of lung and bladder cancer mortality in Chile related to arsenic in drinking water. *J Natl Cancer Inst* 2007;99:920–8. doi:10.1093/jnci/djm004 PMID:17565158
- Chang C-C, Ho S-C, Tsai S-S, Yang C-Y. Ischemic heart disease mortality reduction in an arseniasis-endemic area in southwestern Taiwan after a switch in the tap-water supply system. *J Toxicol Environ Health A* 2004;67:1353–61. doi:10.1080/15287390490471451 PMID:15371236