Manuscript Number: LITHOS3624R1

Title: Highly depleted cratonic mantle in West Greenland extending into diamond stability field in the Proterozoic

Article Type: Research Paper

Keywords: Greenland; Qeqertaa; SCLM; North Atlantic craton; dunite

Corresponding Author: Mr. Kristoffer Szilas, Ph.D.

Corresponding Author's Institution: Lamont-Doherty Earth Observatory

First Author: Stefan Bernstein, Ph.D.

Order of Authors: Stefan Bernstein, Ph.D.; Kristoffer Szilas, Ph.D.; Peter B Kelemen, Ph.D.

Abstract: Abstract

This study presents electron microprobe data for dunite xenoliths from a lamprophyre dyke located on the island of Qeqertaa, West Greenland. The minimum age of this dyke is Palaeoproterozoic and it experienced amphibolite facies metamorphism and deformation during that era. The samples consist of nearly 200 xenoliths with a size range of 0.5-8 cm. These dunite xenoliths have olivine Mg#, that range from 80.3 to 94.6 ($n = 579$) with a mean of 92.6. Orthopyroxene is found in three xenoliths and garnet in five others. The latter suggests the depth of the Qeqertaa xenolith suite to be near the diamond stability-field, which is substantiated by the finding of diamonds in bulk samples of the Qeqertaa dyke. This further indicates the presence of a lithospheric mantle domain dominated by high-Mg# dunite to this depth in Palaeoproterozoic time. Cr-rich spinel, in the 0.1-0.2 mm size range, is found within and between olivine grains in individual xenoliths. These Cr-spinels yield Fe-Mg exchange temperatures of 400-600°C. However, the presence of intermediate spinel compositions spanning the lower temperature solvus suggests that equilibration temperatures were >550°C. Fe^{3+} , expressed as $100 \times Fe^{3+} / (Fe^{3+} + Al + Cr)$, is shown to be a useful parameter in order to screen for altered spinel ($Fe^{3+} > 10$) with disturbed Mg# and Cr#. The screened spinel data ($Fe^{3+} < 10$) show a distinctly different trend in terms of spinel Cr# versus Mg#, compared to unmetamorphosed xenoliths in Tertiary lavas and dikes from Ubekendt Ejland and Wiedemann Fjord, respectively, also located within the North Atlantic craton. This difference likely reflects amphibolite facies metamorphic resetting of the Qeqertaa xenolith suite by Fe-Mg exchange. Given the similarity of the Qeqertaa xenolith suite with the Ubekendt and Wiedemann suites, in terms of their olivine Mg# and spinel Cr# distribution, high-Mg# dunite is likely to be an important component of the subcontinental lithospheric mantle beneath the North Atlantic craton and appears to have spanned a vertical distance of at least 150 km in this region, even during the Palaeoproterozoic.

Abstract

This study presents electron microprobe data for dunite xenoliths from a lamprophyre dyke located on the island of Qeqertaa, West Greenland. The minimum age of this dyke is Palaeoproterozoic and it experienced amphibolite facies metamorphism and deformation during that era. The samples consist of nearly 200 xenoliths with a size range of 0.5-8 cm. These dunite xenoliths have olivine Mg#, that range from 80.3 to 94.6 ($n = 579$) with a mean of 92.6. Orthopyroxene is found in three xenoliths and garnet in five others. The latter suggests the depth of the Qeqertaa xenolith suite to be near the diamond stability-field, which is substantiated by the finding of diamonds in bulk samples of the Qeqertaa dyke. This further indicates the presence of a lithospheric mantle domain dominated by high-Mg# dunite to this depth in Palaeoproterozoic time. Cr-rich spinel, in the 0.1-0.2 mm size range, is found within and between olivine grains in individual xenoliths. These Cr-spinels yield Fe-Mg exchange temperatures of 400-600°C. However, the presence of intermediate spinel compositions spanning the lower temperature solvus suggests that equilibration temperatures were >550°C. $Fe^{3+}\#$, expressed as $100 \times Fe^{3+} / (Fe^{3+} + Al + Cr)$, is shown to be a useful parameter in order to screen for altered spinel ($Fe^{3+}\# > 10$) with disturbed Mg# and Cr#. The screened spinel data ($Fe^{3+}\# < 10$) show a distinctly different trend in terms of spinel Cr# versus Mg#, compared to unmetamorphosed xenoliths in Tertiary lavas and dikes from Ubekendt Ejland and Wiedemann Fjord, respectively, also located within the North Atlantic craton. This difference likely reflects amphibolite facies metamorphic resetting of the Qeqertaa xenolith suite by Fe-Mg exchange. Given the similarity of the Qeqertaa xenolith suite with the Ubekendt and Wiedemann suites, in terms of their olivine Mg# and spinel Cr# distribution, high-Mg# dunite is likely to be an important component of the subcontinental lithospheric mantle beneath the North Atlantic craton and appears to have spanned a vertical distance of at least 150 km in this region, even during the Palaeoproterozoic.

- We present EMP data for a new suite of dunite xenoliths from a lamprophyre dyke in West Greenland.
- Olivine Mg# averages 92.6 and garnet is present together with macrodiamonds.
- Spinel Cr# for this suite appears overprinted by metamorphism and formation of mica coatings.

Highly depleted cratonic mantle in West Greenland extending into diamond stability field in the Proterozoic

Stefan Bernstein ^a, Kristoffer Szilas ^{b, *}, Peter B. Kelemen ^b

^a Avannaa Resources, Dronningens Tværgade 48 st.tv, 1302 Copenhagen K, DK

^b Lamont-Doherty Earth Observatory, PO Box 1000, Palisades, NY 10964-8000, USA

* Corresponding author.

E-mail: kszilas@ldeo.columbia.edu (K. Szilas)

Abstract

This study presents electron microprobe data for dunite xenoliths from a lamprophyre dyke located on the island of Qeqertaa, West Greenland. The minimum age of this dyke is Palaeoproterozoic and it experienced amphibolite facies metamorphism and deformation during that era. The samples consist of nearly 200 xenoliths with a size range of 0.5-8 cm. These dunite xenoliths have olivine Mg#, that range from 80.3 to 94.6 (n = 579) with a mean of 92.6. Orthopyroxene is found in three xenoliths and garnet in five others. The latter suggests the depth of the Qeqertaa xenolith suite to be near the diamond stability-field, which is substantiated by the finding of diamonds in bulk samples of the Qeqertaa dyke. This further indicates the presence of a lithospheric mantle domain dominated by high-Mg# dunite to this depth in Palaeoproterozoic time. Cr-rich spinel, in the 0.1-0.2 mm size range, is found within and between olivine grains in individual xenoliths. These Cr-spinels yield Fe-Mg exchange temperatures of 400-600°C. However, the presence of intermediate spinel compositions spanning the lower temperature solvus suggests that equilibration temperatures were >550°C. $\text{Fe}^{3+}\#$, expressed as $100 \times \text{Fe}^{3+} / (\text{Fe}^{3+} + \text{Al} + \text{Cr})$, is shown to be a useful parameter in order to screen for altered spinel ($\text{Fe}^{3+}\# > 10$) with disturbed Mg# and Cr#. The screened spinel data ($\text{Fe}^{3+}\# < 10$) show a distinctly different trend in terms of spinel

Cr# versus Mg#, compared to unmetamorphosed xenoliths in Tertiary lavas and dikes from Ubekendt Ejland and Wiedemann Fjord, respectively, also located within the North Atlantic craton. This difference likely reflects amphibolite facies metamorphic resetting of the Qeqertaa xenolith suite by Fe-Mg exchange. Given the similarity of the Qeqertaa xenolith suite with the Ubekendt and Wiedemann suites, in terms of their olivine Mg# and spinel Cr# distribution, high-Mg# dunite is likely to be an important component of the subcontinental lithospheric mantle beneath the North Atlantic craton and appears to have spanned a vertical distance of at least 150 km in this region, even during the Palaeoproterozoic.

Keywords: Greenland; Qeqertaa; SCLM; North Atlantic craton; dunite

1. Introduction

Determining the composition of the subcontinental lithospheric mantle (SCLM) has implications for our understanding of the crust-mantle system and its evolution through time. Previous studies on SCLM xenoliths from Greenland have shown the occurrence of nearly monomineralic dunites consisting of remarkably refractory olivine with molar $\text{Mg}/(\text{Mg}+\text{Fe}^{2+})$, or Mg#, averaging about 92.8 (e.g. Bernstein et al., 1998, 2006, Bizzarro and Stevenson, 2003; Garrit, 2000; Wittig et al., 2008). The Qeqertaa xenolith suite presented in this study shows equally refractory olivine compositions. This growing body of data on cratonic mantle xenoliths from Greenland suggests that such olivine-rich mantle is common here, and perhaps comprises a large proportion of the lithospheric mantle beneath substantial parts of Greenland.

High and consistent Mg# in olivine is thought to reflect partial melting of the mantle to the point of exhaustion of orthopyroxene (Bernstein et al., 2007). The implied high degree of melting (37-45%; Bernstein, 1998; Herzberg, 2004) is not achieved in any current geological environment and is thus thought to reflect a hotter mantle during the formation of cratonic SCLM. This is in

52 agreement with the generally Archaean Re-depletion ages of cratonic SCLM xenoliths (e.g.
53 Hanghøj et al., 2001; Pearson et al., 2003; Shirey and Walker, 1998; Wittig, 2010) and the inferred
54 hotter mantle at that time (Herzberg et al., 2010). However, the exact formation environment is still
55 debated with one model proposing a single-stage process in a polybaric melting column either in a
56 spreading ridge or plume environment (e.g. Aulbach et al., 2011; Bernstein et al., 1998, 2006;
57 Griffin et al., 2009; Herzberg et al., 2010; Kelemen et al. 1998), whereas another model proposes
58 flux melting of previously depleted harzburgite in a subduction zone setting (e.g. Canil, 2004; Lee,
59 2006; Wittig et al., 2008).

60 In addition to documenting the composition of the Palaeoproterozoic xenolith suite at Qeqertaa
61 this study also shows that although parameters such as Mg#, and molar Cr/(Cr+Al), or Cr#, may at
62 first appear to retain information of the primary composition of a mantle xenolith suite, examination
63 of molar $\text{Fe}^{3+}/(\text{Cr}+\text{Al}+\text{Fe}^{3+})$, or $\text{Fe}^{3+}\#$, in associated spinels reveals a history of alteration that
64 strongly modified both Mg# and Cr# so that even visually unaltered spinel had its chemistry
65 overprinted during amphibolite facies metamorphism. Thus, spinel compositions, which are widely
66 thought to be reliable indicators of primary igneous conditions in metamorphic intrusive rocks (e.g.
67 Barnes, 2000) may reflect re-equilibration during metamorphic events (Evans and Frost, 1975; Sack
68 and Ghiorso 1991).

69

70 2. Geology

71

72 The Qeqertaa xenolith suite is hosted by an up to 6 m wide vertical dyke of ultramafic
73 lamprophyric affinity, which crops out on the small island Qeqertaa, some 50 km north of Ilulissat
74 at 69°38N, 50°38W (Fig. 1). The dyke is one of many that cut late Archaean gneisses and
75 supracrustal rocks in the Ataa area, eastern Disko Bay (Garde and Steenfelt, 1999; Larsen and Rex,
76 1992; Marker and Knudsen, 1989). The Qeqertaa dyke has not been dated, but similar intrusions in
77 the Ataa area have yielded K-Ar ages of 1782 Ma and 1743 Ma, both ± 70 Ma (Larsen and Rex,

1992). Because the dykes have been affected both by metamorphism and deformation linked to the Palaeoproterozoic Rinkian-Nagsoqtuqidian orogeny, these ages could represent metamorphic overprinting and thus represent minimum ages, as pointed out by Larsen and Rex (1992). Structural interpretation suggests that the Ataa region can be divided into a series of crustal blocks with distinct tectono-magmatic history (Garde and Steenfelt, 1999) and the Qeqertaa dyke is situated in the border zone between the Ataa domain to the north and the Rodebay domain to the south.

The Qeqertaa xenoliths are rounded to subangular, with a size range of 0.5 cm to 8 cm in the longest dimension. In several places along the dyke, the xenoliths are so abundant as to make a clast-supported network. The dyke has brecciated contacts with tonalitic gneisses, and is deformed with pinch and swell structures. Apophyses are often sheared into tight isoclinal folds. Deformation mainly affected the matrix, which often shows carbonate crystallization in pressure shadows of the xenoliths and matrix foliation wrapping around individual xenoliths (Fig. 2). Matrix mineralogy is dominated by tremolite, mica, carbonate, ilmenite and iron oxides. Of these minerals, only mica and ilmenite are thought to remain from the primary mineralogy of the dyke matrix, although even these minerals show evidence of alteration and recrystallization, manifested as oxide exsolution along cleavage planes in mica, and exsolution lamellae and oxidized microcrystalline overgrowth zones on ilmenite.

A collection of nearly 200 xenolith samples form the basis of this study. Individual xenoliths larger than about 3 cm were cut into several slices 5-8 mm thick and all xenoliths were inspected visually before a subset of 119 xenoliths of varying size were prepared for standard polished thin sections and analysed by electron microprobe. The xenolith suite as a whole appeared very homogeneous in terms of mineral mode, texture and grain size. The xenoliths are all dunites with only a few samples containing spinel, garnet, mica or orthopyroxene (see below). A few xenoliths are completely serpentinised, while others are relatively fresh peridotite with only local alteration along minor cracks and veins. However, as shown below, all studied xenoliths have experienced some degree of chemical modification of their primary minerals even though they at a first glance

104 appear nearly unaltered. Typically, in standard thin sections olivine grain margins are yellowish to
105 light brown, in some samples more dusty brown (Figs. 2 and 3). Such margins can be relatively
106 wide (1-2 mm), but are mostly in the order of 0.2-0.6 mm, such as in the example of xenolith
107 sample qq-2 in Fig. 3. Spinel is often oxidized, with irregular grain margins and is opaque in
108 standard thin sections. In rare cases, the spinel has retained a brownish translucent core. Secondary
109 oxide, mainly magnetite, is common along veins and cracks within olivine grains and at the rim of
110 individual xenoliths (Fig. 3). Some examples of xenoliths and their textures are given in Fig. 2a-d.

111 All xenoliths are coarse protogranular, following the terminology of Mercier and Nicolas (1975).
112 Grain size varies from 1-2 mm to >30 mm for olivine. Representative smaller xenoliths are
113 composed of only a few olivine grains. Well-rounded olivine megacrysts up to 30 mm are
114 frequently found in the matrix. An example of one coarse grained xenolith is given in Fig. 3, with
115 olivine grain size ranging 1-15 mm. Orthopyroxene, found in three xenoliths, occurs as ~ 1 mm
116 wide circular inclusions in large olivine grains (Fig. 2). Garnet, found in five xenoliths (out of 200),
117 occurs as rounded 1-2 mm grains, but one 4 mm rounded garnet grain appears in xenolith #463737-
118 1. Only one garnet is partly preserved (sample #463711), while all other grains have been
119 completely replaced by kelyphite (Fig. 2c). Cr-rich spinel is found as minute anhedral inclusions in
120 the 0.1-0.2 mm size range within olivine, or between olivine grains. Relic brown translucent spinel
121 is sometimes found, jacketed by opaque oxide, and nearly always with a thin (20-50 μm wide)
122 coating of mica forming the outer contact to host olivine (Figs. 6a, 7a, 8a). Chrome-bearing
123 magnetite occurs throughout the xenoliths, as irregular larger grains (1-2 mm), possibly replacing
124 primary chrome spinel. It also occurs as smaller individual grains or aggregates of smaller grains
125 (0.1-0.3 mm), often associated with fractures or veins in olivine. These chrome bearing magnetite
126 grains likely represent alteration products. Primary mica is found in two xenoliths. In sample
127 #463707-2, a 2 mm long colourless mica grain is enclosed by olivine. In sample #463706-3 similar
128 colourless mica is found interstitially to medium grained olivine crystals.

129 Several xenoliths show evidence of fracturing, and the presence of carbonate-magnetite
130 intergrowths in such fractures suggests that the xenoliths became disaggregated within the
131 lamprophyre dyke during crustal deformation. This can be appreciated in Fig. 2a, showing
132 secondary carbonate mineralization along extensive fracturing of olivine in xenolith #1 in sample
133 #463715 connected to similar carbonate-Fe-oxide intergrowth in the matrix. Another example of
134 this is presented in Fig. 2b (sample #463728), where carbonate-Fe-oxide intergrowth forms 'horns'
135 at the edges of xenoliths #1 and #2 - probably representing growth in pressure shadows.

136 Tremolite and talc is coexisting with olivine in some of the altered portions of the xenoliths and
137 provides evidence for premetamorphic alteration of at least parts of the olivine grains. A collection
138 of micro- and macrodiamonds have been recovered from several bulk samples of the Qeqertaa dyke
139 (Marmo et al., 2012) during exploration activities from 2007-2012.

140

141 **3. Analytical technique**

142

143 Minerals grains were analysed on standard 40 µm polished thin sections, using the JEOL
144 electron microprobe at the Institute of Geology and Geography, University of Copenhagen. All
145 elements in silicates were measured by WDS, with 20 s peak count time for Na, Mg, Fe, Si, and Ti,
146 while Cr, Ni, Ca and Al were measured with 40 s peak count time. Natural and synthetic standards
147 were measured at the beginning and at the end of each session. For traverses in spinel, an analytical
148 routine optimized for oxides was utilized and the elements analysed were Mg, Fe, Si, Ca, Ti, Cr,
149 Mn, Fe, Ni and Zn. Peak count time for Zn was 40 s and 10 s for remaining elements. Background
150 count time on either side of the peak was half that of peak count time for both silicates and spinels.
151 Table 1 in the online supplementary material lists a series of representative mineral data. The
152 mineral data presented there represents the core compositions, typically 2-3 analyses, and for
153 orthopyroxene and garnet, an average of 4-6 analyses.

154

155 4. Results

156

157 4.1. Olivine chemistry

158

159 There is a large compositional variation in olivine. The overall range in Mg#
160 ($100 \times \text{Mg}/(\text{Mg} + \text{Fe}^{2+})$) out of almost 600 analyses is from 80.3 to 94.6 as shown in the insert in Fig.
161 4. Ni varies from 1328 ppm to 4008 ppm (median value of 2727 ppm) and does not correlate with
162 Mg#. Because olivine composition cannot be related to the textural state of olivine, i.e. xenocrystic
163 or xenolithic (not shown), all olivine in the following is referred to as xenoliths. Fig. 4 also shows
164 the distribution of Mg# in olivine in 119 individual xenoliths, as presented in Table 1 (see online
165 supplementary material). While this data set represents core compositions, the range in Mg# is
166 similarly large, spanning Mg# 80-94. Neither of the data sets (all olivine analyses and xenolith
167 cores only) show a good correlation of Ni versus Mg# (see Fig. 4, insert). Several olivine grains
168 have been analysed along traverses from crystal rim to core. The data are presented in Fig. 5 in
169 terms of the variations in Mg# and Ni contents. Fig. 5a shows a traverse over about 0.3 mm in the
170 core of a large olivine crystal. The variation in Mg# is moderate in the range 93.4-94.1, while Ni is
171 in the range 2295-2955 ppm. These ranges are only slightly larger than what is known from olivine
172 in fresh, unaltered mantle xenoliths from Tertiary volcanics and dykes elsewhere in Greenland
173 (Bernstein et al., 1998, 2006).

174 Some Qeqertaa xenoliths have olivine crystals, such as samples #qq1 and #qq4 (Fig. 5b and 5c),
175 that show similarly consistent Mg# and Ni concentrations over more than one millimetre traverse
176 and with only a modest zoning with small decreases in Mg# over the outer 0.1 mm of the crystal.
177 These crystals have only weakly altered olivine rims (#qq1, Fig. 5b) or no detectable alteration
178 (#qq4-olivine2, Fig. 5c). Other samples contain olivine grains, which show more pronounced
179 zoning, such as #qq2a-olivine1 and #qq3-olivine1 (Fig. 5d and 5e), with core compositions of 93.4
180 and 92.0, respectively, and rim compositions of 90.3 and 87.7. Ni varies from an average value of

181 about 2500 ppm in the high Mg# region, decreasing to 2000 and then to 1600 ppm at the grain
182 edges. These two olivine crystals are also typical of the more altered versions of olivine, with thick
183 0.2-0.3 mm alteration rims marked by brown coloration (Fig. 3a). The colour seems to stem from
184 minute oxide inclusions, typically less than 1 μm , in the olivine, mainly Fe-oxide, as judged from
185 microprobe data.

186 One olivine crystal was analysed with a traverse (Fig. 5f) that runs from crystal core to the edge
187 against a spinel grain (Fig. 7a). The olivine crystal is homogeneous apart from one fracture, which
188 coincides with a small excursion from a steady level at around Mg# 92.9 to 92.2, accompanied by a
189 small decrease in Ni. The olivine grain terminates at a thin veneer (20 μm wide) of mica that
190 surrounds the spinel grain (Fig. 7a), and the olivine attains its high Mg# and Ni content approaching
191 this contact (Fig. 5f).

192

193 4.2. *Spinel chemistry*

194

195 As discussed above, irregular spinel grains or aggregates of magnetite and chromian magnetite
196 are found along cracks and veins in the olivine xenoliths. Spinel textures and occurrence suggests
197 that these irregular grains and aggregates are of secondary origin. Spinel grains with similar texture
198 and composition are also ubiquitous in the matrix and will not be discussed further.

199 Spinel of likely primary origin is found in portions of the xenoliths, where alteration has been
200 less intense. Some spinel grains interpreted as relic primary spinel are present in the centre of
201 irregular masses or aggregates of iron oxide grains. On the basis of microscopy and back-scatter
202 electronic images, it is apparent that two basic types of relic, primary spinel grains occur: 1) some
203 (rare) with recognizable primary grains boundaries, only narrow alteration rims and a less modified
204 core, and 2) more intensely altered grains, with obscured primary grain boundaries, often with
205 overgrowths of chromian magnetite.

206 The best preserved spinel grains are found as two small (0.1-0.3 mm) slightly translucent grains
207 completely enclosed in a large (12 mm) olivine grain (Fig. 6a). The compositional variation of the
208 core of this olivine grain is presented in Fig. 5a. The two spinel grains are compositionally similar
209 with Cr#, calculated as $Cr\# = (100 \times Cr / (Cr + Al))$, in the spinel cores of 60.0 ± 0.5 , increasing to a
210 maximum of about 62 at grain edges (Fig. 6b). Mg# and Al#, calculated as $Al\# =$
211 $(100 \times Al / (Fe^{3+} + Al + Cr))$, also show restricted ranges at around 56 and 39, respectively, decreasing
212 slightly at grain edges. $Fe^{3+}\#$ is very low, mostly between 1 and 2, and increasing slightly to 3-4 at
213 grain edges. ZnO shows slight variation around 0.5 wt.%, while both NiO and TiO_2 is at or below
214 detection limit (Fig. 6d). As examples of the relatively well-preserved spinels of type 1), but with
215 pronounced alteration, Figs. 7a and 8a show two spinel grains in the 100 μm size range. Texturally
216 the two grains are similar, with a translucent core, and a pitted and opaque rim. Spinel grain #qq4b-
217 sp1 (Fig. 7a) is rounded and anhedral, typical for spinel grains in xenoliths of cratonic mantle
218 peridotite, while grain #qq3-sp1 (Fig. 8a) is subhedral. Fine-grained mica aggregates separate the
219 spinel from hosting olivine grains. Chemically, the two spinel grains share some characteristics with
220 relatively constant levels of Cr#, Mg#, $Fe^{3+}\#$, and Al# in the core, that change markedly at the grain
221 edges towards elevated Cr# and $Fe^{3+}\#$ and lower Mg# and Al#. This zoning is also pronounced for
222 ZnO, which varies from about 1 wt.% in the centre of the spinel grains to 3 wt.% towards the rim
223 for #qq4-sp1 (Fig. 7d). For spinel #qq3-sp1, the variation and absolute concentrations in ZnO is less
224 extreme with about 0.4-0.7 wt.% (Fig. 8d). For titanium and nickel, core concentrations are ~ 0.1
225 wt.% increasing to 0.5 wt.% (TiO_2) and ~ 0.3 wt.% (NiO) at the crystal rim (Fig. 7d). Spinel #qq3-
226 sp1 again shows less variation, with 0.2-0.3 wt.% TiO_2 and ≤ 0.1 wt.% NiO (Fig. 8d). MnO follows
227 the pattern of ZnO at similar concentration levels (not shown) for all analysed spinel grains.

228 As an example of spinels of type 2), which are more altered and anhedral, Fig. 9 illustrates one
229 grain from sample #qq4, with a short analytical traverse from grain boundary to some way into the
230 core. As for the type 1 spinels in Figs. 7 and 8, a mosaic of mica crystals completely surrounds this
231 spinel, separating it from the host olivine. In this case the mica zone is 50-100 μm wide. The spinel

232 is opaque and the backscatter electron image (Fig. 9a) does not reveal significant zoning, although
233 the rounded shape of the primary spinel can be detected inside the ragged rim. In contrast to the
234 type 1 spinel grains, type 2 shows little variation from core to rim, and instead has extreme Cr#,
235 approaching 100. Mg# is low (<20) and Fe³⁺# high (>50). Concentrations of minor elements ZnO
236 and TiO₂ are considerably less variable than for type 1 spinel, and typically with values at 0.7-1.0
237 wt.%, while NiO is low between 0.1 and 0.2 wt.% (Fig. 9d).

238 From the data presented above it is clear that all spinel grains have undergone variable degrees
239 of chemical modification since the lamprophyre hosting the xenoliths was emplaced. Only a few
240 spinel grains have been analysed with electron microprobe traverses, so it is important to use a
241 chemical parameter that can distinguish between highly altered spinel cores and the less altered
242 cores that potentially retain their primary Cr#. One such parameter is Fe³⁺#, which varies from 1.3
243 to 98.0 in Qeqertaa samples. This is in marked contrast to unaltered and unzoned spinel in mantle
244 xenoliths in Tertiary lavas and dykes in West and East Greenland, which all have Fe³⁺# <10 in
245 spinels that span a Cr# range of 25-95 (Fig. 10). By contrast, Table 1 in the online supplementary
246 material lists all spinel data from Qeqertaa that have Fe³⁺# <10; only 20 samples have such low
247 values.

248 Fig. 11 illustrates the compositions of Qeqertaa spinel with Fe³⁺# <10. The markedly lower Mg#
249 at a given Cr#, compared to the xenolith suites of Ubekendt Ejland and Wiedemann Fjord, is an
250 indication of lower temperature Fe-Mg exchange with olivine and other silicates in the Qeqertaa
251 spinels. The Qeqertaa samples yield olivine-spinel Fe-Mg exchange temperatures of 400-600°C
252 (Ballhaus et al., 1991; Sack and Ghiorso, 1991), which is substantially lower than Ubekendt Ejland
253 (833-960°C; Bernstein et al., 2006) and Wiedemann Fjord (650-995°C; Bernstein et al., 1998).

254

255 4.3. Orthopyroxene chemistry

256

257 The three orthopyroxene grains from three individual xenoliths show a restricted variation, with
258 Mg# spanning 92.7-93.7, and with Cr₂O₃ and Al₂O₃ levels at 0.41-0.52 wt.% and 0.79-0.85 wt.%,
259 respectively (Table 1, online supplementary material). CaO is ca. 1.1 wt.% for two samples, while
260 the remaining sample has only 0.3 wt.% CaO. The three dunite xenoliths that contain orthopyroxene
261 have no garnet, but the high Cr# (26-30) of all three orthopyroxene grains suggests that they have
262 equilibrated with garnet, as orthopyroxene from garnet-free peridotite xenoliths have significantly
263 lower Cr#, mostly less than 20 (Fig. 12). An important note here is that the distinction between
264 orthopyroxene from garnet-bearing and garnet-free peridotite disappears for orthopyroxene with
265 extremely low contents of Al₂O₃ (<0.2 wt.%), that have variable Cr#, from 4-96 (Fig. 12). This is
266 possibly an artefact of poor analytical accuracy or extreme Al depletion during mantle melting and
267 melt extraction in spinel stability field (Bernstein et al., 2007; Stachel et al., 1998). However, the
268 distinction appears robust for orthopyroxene with Al₂O₃ exceeding 0.2 wt.%.

269 Because of the lack of coexisting phases, such as garnet or clinopyroxene, it is not possible to
270 calculate equilibrium temperatures for the garnet-bearing assemblages, and it is indeed questionable
271 if such calculations would be meaningful, given that the xenoliths were subjected to amphibolite
272 facies metamorphism after emplacement of the Qeqertaa dyke.

273

274 **5. Discussion**

275

276 *5.1. Metamorphic overprinting*

277

278 The strongly zoned nature of olivine and chrome spinel from the xenoliths of Qeqertaa is a
279 typical aspect of prograde metamorphism of altered peridotite (e.g. Evans and Frost, 1975; Nozaka,
280 2003; Trommsdorff et al., 1998; Vance and Dungan, 1977). The Qeqertaa olivines with brown,
281 inclusion-rich grain margins show textural similarity to hydrothermally altered olivine xenocrysts
282 and xenoliths from Venetia kimberlites, South Africa (Stripp et al., 2006), where thin talc rims form

on serpentinised olivine grains. In a study of olivine compositional variation in Chugoku peridotite (Japan), which has been subjected to steep local metamorphic gradients, Nozaka (2003) demonstrates how primary olivine with Mg# around 91 can attain both higher (to 97.6) and lower (to 86) Mg# during prograde reactions. The assemblage at Qeqertaa of coexisting olivine, tremolite and talc is analogous to metamorphic zone 3 of Nozaka, (2003, and references therein), where essentially all olivine is metamorphic. The metamorphic olivine in Chugoku peridotites forms overgrowths, along distinct grain boundaries and 'healed' cracks or veins, where olivine replaces serpentine. Magnetite-free serpentine is thought to be the precursor of high-Mg# olivine, while low-Mg# olivine is thought to crystallize from magnetite-bearing serpentine. The olivine in Qeqertaa xenoliths always shows normal zoning with low-Mg# zones crowded with minute Fe-oxide inclusions. These low-Mg# rims are thus likely to represent recrystallized, partly serpentinised olivine with Fe-oxide grains remaining from the hydrothermally reacted olivine.

In Qeqertaa xenoliths, all olivine grains show some reaction along grain boundaries, evident from the presence of either hydrous phases, such as talc, or fine-grained Fe-oxide inclusions. However, many olivine grain centres are colourless, transparent and inclusion-free down to the resolution of the electron microprobe. These olivine cores coincide with maximum Mg# in the individual olivine and suggest that the cores represent primary olivine that escaped extensive serpentinisation. Likewise, the typical coarse protogranular texture (Fig. 3) suggests that the Qeqertaa xenoliths are not entirely recrystallized from serpentine or talc during prograde metamorphism, as such recrystallization commonly forms equigranular olivine aggregates or highly elongate olivine resembling spinifex texture (e.g. Trommsdorff et al., 1998). Another piece of evidence for the colourless olivine to have escaped the initial serpentinization is the presence of inclusions of pale, inclusion-free phlogopite crystals that are in strong contrast to the dark-brown, inclusion-rich mica phenocrysts or xenocrysts in the groundmass.

The relatively robust nature of chromite makes these a potential powerful recorder of alteration processes (e.g. Barnes, 2000; Evans and Frost, 1975; Kimball, 1990). The types of chemical zoning

309 depicted in Figs. 6-9 can be explained in terms of hydrothermal alteration followed by prograde
310 metamorphism from primary chromite with slightly altered crystal rims to magnetite rimmed spinel
311 grains with equilibrated cores. The high ZnO (mostly at 0.4-3.0 wt.%) present in all spinel grains
312 with electron probe traverses, show that even spinel cores have been modified by metamorphic
313 reactions, because primary spinels have ZnO at the 0.1 wt.% level, at least in komatiite flows
314 (Barnes, 2000). The exception is the spinel portrayed in Fig. 6a, which perhaps escaped strong
315 modification. Certainly, some spinel has been altered to magnetite, which is found in cracks and
316 veins, as well as along olivine grain boundaries (Fig. 3). This suggests at least some degree of early
317 partial serpentinisation and later recrystallization of olivine, which would then leave magnetite
318 interstitially between olivine crystals as seen in Fig. 3.

319 The spinel grains in 463722 and #qq-4b (Figs. 6 and 7) are morphologically the least altered
320 spinel grains of those analysed. The lack of a thick, euhedral Fe-spinel jacket suggests that these
321 grains have remained relatively undisturbed during metamorphism. While the zoning in the two
322 spinel grains in 463722 is very modest, #qq-4b shows a more pronounced decrease of magnesium
323 and aluminium towards the rim and around the transecting crack, which is typical for chromite
324 equilibrating with olivine, talc, or ferroan magnesite (for Mg) and chlorite (for Al) (Barnes, 2000).
325 In the case of the Qeqertaa xenoliths, the samples lack chlorite. However, Cr-Al exchange between
326 spinel and mica, as in the following simplified reaction, would decrease the Al content of spinel:

327

329

330 Instead or in addition, metamorphic reactions forming mica from talc + olivine + aluminous
331 spinel + fluid, as in the following simplified example, could also explain the high Cr# in remaining
332 Qeqertaa spinels:

333

336

337

338 The occurrence of one or both of these reactions is consistent with the phlogopite-rich
 339 composition of the mica that coats the spinel in for example Fig. 7a) at the end of line-1 (see
 340 supplementary data Table 1).

341 Following Barnes' (2000) study, serpentinisation of dunite xenoliths and olivine in the
 342 lamprophyre matrix results in decomposition of olivine and release of Zn. Chromite absorbs Zn,
 343 resulting in a steep concentration profile from high ZnO at altered chromite rims to very low
 344 concentration (<0.1 wt.%) at unaltered chromite cores. Prograde metamorphism to amphibolite
 345 facies subsequently redistributes Zn in the chromite grains and tends to homogenize concentration
 346 differences, which in turn also results in elevated Zn in grains that have retained their primary Cr#
 347 (Barnes, 2000).

348 In a more extreme case, such as in Fig. 9 (#qq4, line4) aluminium is completely lost, along with
 349 most of the magnesium. The overgrowth of Fe-rich spinel exhibits crystal faces, but still contains
 350 appreciable amounts of Cr. In fact, only a few of the analysed chrome spinel grains have pure
 351 magnetite rims. In a plot of all spinel analyses in terms of their trivalent cations (Fig. 13) these
 352 highly altered grains (as in Fig. 9), and altered spinel rims (Figs. 7-8) plot along the Cr-Fe³⁺ join,
 353 clustering at Cr₃₀Fe³⁺₇₀-Cr₅₀Fe³⁺₅₀ and connecting to the main body of spinel data along the Al-Cr
 354 join at low (generally <10%) Fe³⁺. These intermediate spinel compositions suggest extensive solid
 355 solution between the spinel end-member components (Fe,Mg)Cr₂O₄, (Fe,Mg)Al₂O₄ and
 356 (Fe,Mg)Fe₂O₄ and thus equilibration temperatures above 600°C according to Sack and Ghiorso
 357 (1991). At temperatures at or below 550°C the solid solution stability field is significantly reduced
 358 by miscibility gaps, which expand with decreasing temperature, resulting in formation of coexisting

359 chromite and Cr-poor magnetite (see also Barnes, 2000). The presence of intermediate
360 compositions in the Qeqertaa xenoliths suggests crystallization above the solvus

361 Fig. 10 shows that most spinel analyses have high $\text{Fe}^{3+}\#$ (>10) reflecting the effects of
362 hydrothermal alteration and/or amphibolite facies metamorphism. However, even spinel with $\text{Fe}^{3+}\#$
363 <10 have lost their primary Mg# as seen in Fig. 11. The least altered Qeqertaa xenoliths plot along
364 the metamorphic trend identified by Frost and Evans (1975). As in most alpine peridotite massifs,
365 Fe-Mg exchange between spinel and olivine has reduced spinel Mg#.

366 Primary spinel Cr# can in some cases be preserved in metamorphosed peridotites. The range of
367 Cr# in Qeqertaa spinels overlaps with that of Ubekendt Ejland and Wiedemann Fjord xenolith
368 suites, raising the possibility that some crystals retain primary Cr#'s. However, Qeqertaa xenolith
369 spinels plot along the metamorphic Cr# versus Mg# trend of Evans and Frost (1975) for spinels in
370 chlorite-bearing metaperidotites. Very high Cr# and low Mg# in metamorphic spinels were also
371 observed by Barnes and Roeder (2001), further suggesting that Qeqertaa spinel Cr#'s may have
372 been modified by metamorphic reactions, perhaps to some degree even for spinels with low $\text{Fe}^{3+}\#$.

373 In summary, the combined evidence from olivine and spinel points to a history of partial
374 replacement of olivine by serpentine during hydrothermal alteration, followed by prograde
375 metamorphism to amphibolite facies, as suggested by the presence of recrystallized olivine rims, the
376 character of zoned chrome spinel, and olivine-spinel Fe-Mg thermometry.

377

378

379 5.2. Primary olivine compositions

380

381 Considering olivine cores only, the Qeqertaa xenoliths have an average Mg# of 92.6 with a
382 median Mg# of 92.8. Given the considerable zoning, these values are probably minimum estimates
383 of the average and median, values of primary Mg#, because (a) some apparent olivine cores may be
384 from olivine in which the plane of the thin section is parallel to and close to one edge of the crystal,

385 and therefore have broad areas of low-Mg#, and (b) even some of the larger, clear grains may be
386 neoblasts, crystallized from completely serpentinised primary olivine. In any case, olivines from
387 Qeqertaa xenoliths have highly refractory compositions that are typical for dunite and harzburgite
388 xenolith suites from the lithospheric mantle beneath Archaean cratons (e.g. Bernstein et al., 1998,
389 2006, 2007; Boyd, 1989; Boyd et al., 1997; Canil, 2004; Lee and Rudnick, 1999; Menzies, 1990;
390 Pearson et al. 2003; Wittig, 2008).

391 The large range in olivine Ni concentrations (1328 ppm to 4008 ppm) portrayed in Fig. 4 and in
392 detail for a subset of individual olivine crystals in Fig. 5, is not easily explained. Some of the range
393 may reflect the redistribution of Ni during partial serpentinization and subsequent prograde
394 metamorphism as discussed above. One likely example is shown in Fig. 5d, in which the
395 recrystallized olivine rim shows substantially lower Mg# and variable but overall decreasing Ni
396 content compared to olivine core. However, even for colourless, inclusion-free olivine grains or
397 olivine cores, there appears to be considerable variation in Ni content. Our microprobe traverses of
398 seemingly unaltered olivine cores with near-constant Mg# reveal Ni variation spanning 2000-3000
399 ppm over few hundred microns (Fig. 5b, c and f), although one grain (Fig. 5a) shows a much tighter
400 variation in Ni with values between 2400 and 2900 ppm. One potential explanation for such
401 variation could be that all olivine is of metamorphic origin, which in turn implies that the low-
402 temperature hydrothermal alteration altered the entire peridotite mineral assemblage prior to the
403 prograde metamorphism. We find this to be implausible not only on textural grounds as described
404 in section 5.1 but also because rare, large, pale mica crystals in the dunite xenoliths are free of the
405 magnetite inclusions that dominate the altered mica crystals in the groundmass. We find it unlikely
406 that a phlogopite crystal would maintain its textural integrity if it were engulfed in completely
407 serpentinized dunite.

408 During our microprobe work on the olivine, we have not detected any inhomogeneity by back-
409 scatter imagery and the origin of the Ni variation is therefore uncertain. While it could stem from
410 minute sulphide inclusions, we also note that the majority of the analytical values are within the

411 normal range for olivine from olivine-rich, residual mantle peridotites (2000-3500 ppm; e.g.,
412 Bodinier and Godard, 2003). We note that the range in Ni for the Qeqertaa xenolithic olivine is
413 similar to that observed for other xenolith suites in the North Atlantic craton which may suggest
414 that this is a common feature (e.g. Bernstein et al. 1998, 2006; Wittig et al., 2008). One possibility
415 is that early (prior to peridotite entrainment in lamprophyre) melt-rock reaction could be responsible
416 for some of the decrease in Ni content of olivine, such as is observed in dunite replacing harzburgite
417 in the Bay of Island ophiolite (Suhr et al. 2003). However, we do not currently have trace element
418 data to test if melt-rock reactions were significant in this case. Variable Ni concentrations, with very
419 low values in some parts of olivine crystals, could also be due to the presence of sulfides during
420 peridotite recrystallization in the mantle. This is observed in sulfide-rich dunites from Oman
421 (Negishi et al. 2012). However, no sulphides have been observed in any of the Qeqertaa xenoliths.
422 Another potential process that might account for the large variability of Ni in olivine is nickel
423 redistribution during early pervasive serpentinisation, followed by later prograde recrystallisation.
424 We reiterate that the large Ni variation in olivine appears to be a common feature in Greenlandic
425 xenolith suites, regardless of their emplacement age or post-emplacement thermal history. A
426 systematic study of nickel content in olivine for all Greenlandic xenolith suites could address this
427 question.

428 With respect to the nearly monomineralic composition of the Qeqertaa xenoliths, they are similar
429 to the shallow, garnet-free, spinel dunites from Ubekendt Ejland, and the olivine-rich,
430 orthopyroxene-poor spinel harzburgites from Wiedemann Fjord (average ca. 12 wt.%
431 orthopyroxene). All three suites have similar average olivine compositions of Mg# 92.6-92.8
432 (Bernstein et al., 1998, 2006, 2007). The growing body of data on cratonic mantle xenoliths from
433 Greenland suggests that such olivine-rich mantle may be more common here than in other cratons,
434 and perhaps is the most abundant lithology in the shallow, cratonic mantle beneath some parts of
435 Greenland. Sample collections from Southwest Greenland (some 300-800 km south of Disko Bay)
436 have peridotite xenolith suites typically ranging from 75-100% modal olivine, with compositions

437 averaging Mg# 92.5-92.8 (Bizzarro and Stevenson, 2003; Garrit, 2000; Wittig et al., 2008). In other
438 cratons, such as Kaapvaal, Tanzania, Siberia, and Slave, mantle xenolith suites typically have lower
439 modal olivine and higher orthopyroxene, with ranges of 40-80% olivine and 20-60%
440 orthopyroxene, plus additional spinel, clinopyroxene and garnet (see compilations of e.g. Griffin et
441 al., 2003; Herzberg, 1993; Lee, 2006; Pearson et al., 2003).

442 It has long been clear that the high Mg# of olivine in cratonic mantle peridotite exceeds that of
443 olivine from Phanerozoic abyssal peridotite, most orogenic peridotite massifs and most peridotite
444 from arcs (Bonatti and Michael, 1989; Boyd, 1989; Boyd and Mertzman, 1987; Menzies 1990;
445 Nixon and Boyd, 1973) and that this elevated Mg# in olivine and bulk rock compositions likely
446 stems from higher extents of partial melting in the Archaean (Boyd, 1989; Herzberg, 2010; Jordan,
447 1975; O'Hara et al., 1975). The nature of the melting processes has on the other hand been subject
448 to much debate over the last couple of decades, mainly because the orthopyroxene-rich mantle
449 peridotites beneath Kaapvaal and some other cratons, such as Yakutia in Siberia, for many years
450 were considered representative of cratonic mantle. The high proportion of orthopyroxene (often >40
451 wt.%) in these peridotite xenoliths is considered inconsistent with an origin as residues of melt
452 extraction from primitive upper mantle compositions (e.g. Herzberg, 1993; Kelemen et al., 1998;
453 Kesson and Ringwood, 1989; Walter, 1998).

454 Following the discovery of depleted orthopyroxene-poor xenoliths from Greenland and some
455 other cratons (Bernstein et al., 1998, 2007; Boyd and Canil, 1997; Larsen, 1982) it has become
456 widely accepted that such high-Mg#, orthopyroxene poor harzburgites and dunites, and olivine +
457 spinel inclusions in diamonds, can be viewed as archetypical cratonic mantle peridotite (e.g.
458 Bernstein et al., 2007; Pearson and Wittig, 2008) formed by melt extraction of 37-45% (e.g.
459 Herzberg, 2004). The modal composition of orthopyroxene-poor harzburgite and dunite cratonic
460 xenoliths and their high Mg# suggest that they are residues of polybaric decompression melting
461 starting at 4 to 6 GPa and extending to less than 3 GPa (Bernstein et al., 1998, 2006; Herzberg,
462 2004; Kelemen et al., 1998; Walter, 2003). Depending on the nature of melting and melt extraction

463 (“equilibrium porous flow” versus “fractional melting”), this may be consistent with the observation
464 that the trace element composition of SCLM xenoliths from several cratons suggests melting at a
465 shallow depth (< 3 GPa; e.g. Canil, 2004; Wittig et al., 2008). Alternatively, high Mg# and other
466 indications of extreme melt depletion could be the cumulative result of multiple stages, each with a
467 moderate degree of melt extraction. The uniform olivine Mg# of 92.6-92.8 of the cratonic high Mg#
468 dunite may have been controlled by the exhaustion of orthopyroxene during melt extraction,
469 limiting the maximum extent of melt depletion to about 40% in most cases, regardless of the
470 tectonic environment (Bernstein et al., 2007).

471 Likewise, there is a growing consensus that the orthopyroxene-rich nature of mantle peridotite
472 beneath e.g. Kaapvaal, is a result of silica addition through melt-rock reaction, in which silica-rich
473 melts react with a high-Mg# dunite protolith, perhaps in the hanging wall of a subduction zone
474 (Gibson et al., 2008; Griffin et al., 2009; Kelemen et al., 1998; Kesson and Ringwood, 1989; Lee,
475 2006; Pearson and Wittig, 2008; Rudnick et al., 1994; Lee et al., 2011). Indeed, the data and
476 interpretations of Hanghøj et al. (2001) and Bernstein et al. (2007) suggest that Al as well as Si has
477 been added to dunite protoliths in most cratonic harzburgites, and that some cratonic lherzolites are
478 also refertilised dunites, modified by addition of Ca, Fe, Al, Si, and probably many other elements.

479 Pressure and temperature at the time of xenolith entrainment can be calculated for many garnet-
480 bearing peridotite xenoliths. Pressures indicate depths of equilibration in the order of 2.5-7 GPa or
481 about 75-220 km depth (e.g. Lee, 2006). Disregarding xenoliths with clear signs of metasomatic
482 overprinting by a basaltic/basanitic or carbonatitic melt (i.e. presence of abundant diopside,
483 amphibole, mica, carbonate, rutile etc.), there is a surprisingly constant and high Mg# of olivine in
484 cratonic mantle, from 91.5-94.0 for most cratons, with averages spanning 92.0 to 93.0 (e.g. Gaul et
485 al., 2000; Griffin et al., 2003; Pearson and Wittig, 2008), which do not correlate with depth of
486 equilibration. These values are similar to depleted, garnet-free spinel peridotite xenoliths from the
487 cratons, which also have average olivine Mg# of 92.5-92.8 (Bernstein et al., 2007).

488 As noted above, the bulk composition of most garnet-bearing cratonic mantle xenoliths reflects
489 low pressure melting in the absence of garnet (e.g. Canil and Wei, 1992; Kelemen et al., 1998;
490 Stachel et al., 1998) despite their deep residence at the time of entrainment. The trend of bulk
491 compositions, with correlated Ca and heavy rare earth element concentrations, demonstrate that few
492 if any cratonic peridotites contain residual garnet. Instead, most garnets in the cratonic upper mantle
493 must be metamorphic in origin (Kelemen et al., 1998). This interpretation requires that low-pressure
494 peridotite residues of decompression melting have been transported to depth at some later stage,
495 possibly during collision of tectonic plates leading to stacking of the depleted and hence buoyant
496 dunitic restite (e.g. Helmstaedt and Schulze, 1989; Gray and Pysklywec, 2010) or via ascent and
497 accumulation of buoyant, Fe-poor peridotite diapirs (e.g., Oxburgh and Parmentier, 1977, 1978).

498 In this light, the presence of garnet-bearing dunite with average olivine Mg# of 92.6 in the
499 Qeqertaa suite, considered together with the garnet-free Ubekendt and Weidemann xenolith suites,
500 shows that by Palaeoproterozoic time, the mantle beneath this part of the North Atlantic craton
501 included regionally extensive high Mg# dunite extending from depths within the spinel peridotite
502 stability field (Bernstein et al., 2006), deep into the garnet peridotite stability field, and indeed into
503 the diamond stability field as documented by the presence of diamonds in the Qeqertaa dyke
504 (Marmo et al., 2012). Along typical cratonic conductive geotherms (surface heat flow of 40-50
505 mW/m²), this results in a vertical distribution of high-Mg# dunite over 150 km of the lithospheric
506 mantle.

507

508 **6. Conclusions**

509

510 We have shown that yet another SCLM xenolith suite from Greenland records consistently high
511 Mg# in olivine. The Qeqertaa xenoliths have average olivine Mg# of 92.6 and a median value of
512 92.8, which appears to be the dominating composition for SCLM in the North Atlantic craton.

513 The presence of garnet in five xenoliths, suggests an equilibration pressure greater than 2.5 GPa
514 (>75 km depth). The presence of diamonds in the Qeqertaa dyke demonstrates that the SCLM in
515 this region must have been more than 150 km thick prior to the Palaeoproterozoic transport of the
516 xenoliths into the crust.

517 The after emplacement in the crust, Qeqertaa xenoliths likely equilibrated at temperatures
518 slightly above 600°C, as reflected in their spinel compositions, during amphibolite facies
519 metamorphism in the crust. The Cr# versus Mg# trend of spinel from Qeqertaa is different than for
520 xenoliths from Ubekendt Ejland and Wiedemann Fjord (Bernstein et al., 1998, 2006), although they
521 probably shared a similar origin in highly depleted, residual dunites.

522 Unlike spinel Mg#'s, which are typically modified during metamorphism, spinel Cr#'s can in
523 some cases preserve primary, residual compositions. Values of $Fe^{3+}\# < 10$ in some Qeqertaa
524 spinels suggest that primary Cr# may have been preserved in a few cases. However, the Qeqertaa
525 xenolith spinels plot along the metamorphic Cr# versus Mg# trend for chlorite-bearing peridotites
526 identified by Evans and Frost (1975). Our samples do not contain chlorite, but they do contain
527 metamorphic mica separating spinels from host olivine crystals. This suggests that even spinel Cr#
528 can be modified by open behaviour during amphibolite facies metamorphism. Thus, care must be
529 taken when interpreting the geological formation environment of metamorphic mantle rocks, in
530 terms of their spinel composition.

531 The surprisingly monotonous lithology in the 100% dunite Qeqertaa xenolith suite further
532 supports the hypothesis that much of the cratonic mantle is composed of low pressure residues of
533 high degrees of decompression melting, limited by the exhaustion of orthopyroxene (Bernstein et
534 al., 2007). The Qeqertaa data are similar to xenolith data from other areas of the North Atlantic
535 craton, as well as some other cratons as seen in Fig. 14, which supports the hypothesis that olivine-
536 rich dunites and harzburgites with Mg# around 92.8 represent the pristine composition of SCLM,
537 prior to refertilization via melt-rock reaction.

538 **Acknowledgements**

539

540 We thank Jacques Deleuran, Birgit Leer and Marie Leer Jørgensen for assistance during xenolith
541 sampling and Artisk Station (University of Copenhagen) and its ship R/V Porsild for transportation
542 and other field support. Berit Wenzel and Alfons Berger helped with the electron microprobe.
543 Thanks also to Minik Rosing and NordCEE for supporting the senior author. We thank G. Nelson
544 Eby for editorial handling of the manuscript and Claude Herzberg and one anonymous reviewer for
545 constructive comments on an early version.

546

547 **References**

548

549 Arai, S., 1994. Characterization of spinel peridotites by olivine-spinel compositional relationships:
550 Review and interpretation. *Chemical Geology* 113, 191-204.

551

552 Aulbach, S., Stachel, T., Heamann, M., Creaser, R.A., Shirey, S.B., 2011. Formation of cratonic
553 subcontinental lithospheric mantle and complementary komatiite from hybrid plume sources.
554 *Contributions to Mineralogy and Petrology* 161, 947-960.

555

556 Ballhaus, C., Berry, R.F., Green, D.H., 1991. High pressure experimental calibration of the olivine-
557 orthopyroxene-spinel oxygen barometer: implications for the oxidation state of the mantle.
558 *Contributions to Mineralogy and Petrology* 107, 27-40.

559

560 Barnes, S.J., 2000. Chromite in komatiites II. Modification during greenschist to mid-amphibolite
561 facies metamorphism. *Journal of Petrology* 41, 387-409.

562

563 Barnes, S.J., Roeder, P.L., 2001. The range of spinel compositions in terrestrial mafic and
564 ultramafic rocks. *Journal of Petrology* 42, 2279-2302.

565

566 Bernstein, S., Kelemen, P.B., C.K. Brooks, 1998. Depleted spinel harzburgite xenoliths in Tertiary
567 dykes from East Greenland: Restites from high degree melting. *Earth and Planetary Science Letters*
568 154, 221-235.

569

570 Bernstein, S.K., Hanghøj, K., Kelemen, P.B., Brooks, C.K., 2006. Ultra-depleted, shallow cratonic
571 mantle beneath West Greenland: Dunitic xenoliths from Ubekendt Ejland. *Contributions to*
572 *Mineralogy and Petrology* 152, 335-347.

573

574 Bernstein, S., Kelemen, P.B., Hanghøj, K., 2007. Consistent olivine Mg# in cratonic mantle reflects
575 Archean mantle melting to the exhaustion of orthopyroxene. *Geology* 35, 459-462.

576

577 Bizzarro, M., Stevenson, R.K., 2003. Major element composition of the lithospheric mantle under
578 the North Atlantic craton: Evidence from peridotite xenoliths of the Sarfartoq area, southwestern
579 Greenland. *Contributions to Mineralogy and Petrology* 146, 223-240.

580

581 Bodinier, J.-L., Godard, M., 2003. Orogenic, ophiolitic and abyssal peridotites. In: Holland, H.D.,
582 Turekian, K.K. (Eds.), *Treatise on Geochemistry*, vol. 2. Elsevier.

583

584 Bonatti, E., Michael, P.J., 1989. Mantle peridotites from continental rifts to ocean basins to
585 subduction zones. *Earth and Planetary Science Letters* 91, 297-311.

586

587 Boyd, F.R., 1989. Compositional distinction between oceanic and cratonic lithosphere. *Earth and*
588 *Planetary Science Letters* 96, 15-26.

589

590 Boyd, F.R., Mertzman, S.A., 1987. Composition and structure of the Kaapvaal lithosphere, southern
 591 Africa. Magmatic processes. In: B.A. Mysen (Ed.) *Physiochemical Principles*. University Park, PA,
 592 The Geochemical Society, Special Publication 1, 13-24.
 593
 594 Boyd, F.R., Canil, D., 1997. Peridotite xenoliths from the Slave craton, Northwest Territories.
 595 Seventh Annual V. M. Goldschmidt Conference, LPI Contribution No. 921, Lunar and Planetary
 596 Institute, Houston.
 597
 598 Boyd, F.R., Pokhilenko, N.P., Pearson, D.G., Mertzman, S.A., Sobolev, N.V., Finger, L.W., 1997.
 599 Composition of the Siberian cratonic mantle: Evidence from Udachnaya peridotite xenoliths.
 600 *Contributions to Mineralogy and Petrology* 128, 228-246.
 601
 602 Canil, D., 1992. Orthopyroxene stability along the peridotite solidus and the origin of cratonic
 603 lithosphere beneath southern Africa. *Earth and Planetary Science Letters* 111, 83-95.
 604
 605 Canil, D., 2004. Mildly incompatible elements in peridotite and the origins of mantle lithosphere.
 606 *Lithos* 77, 375-393.
 607
 608 Canil, D., Wei, K., 1992. Constraints on the origin of mantle-derived low Ca garnets. *Contributions*
 609 *to Mineralogy and Petrology* 109, 421-430.
 610
 611 Evans, B.W., Frost, B.R., 1975. Chrome-spinel in progressive metamorphism- a preliminary
 612 analysis. *Geochimica et Cosmochimica Acta* 39, 959-972.
 613
 614 Garde, A.A., Steenfelt, A., 1999. Precambrian geology of Nuussuaq and the area north-east of
 615 Disko Bugt, West Greenland. *Geology of Greenland Survey Bulletin* 181, 6-40.

616

617 Garrit, D., 2000. The nature of the Archaean and Proterozoic lithospheric mantle and lower crust in
618 West Greenland illustrated by the geochemistry and petrography of xenoliths from kimberlites.
619 unpublished ph. d. thesis University of Copenhagen, 289.

620

621 Gaul, O.F., Griffin, W.L., O'Reilly, S.Y., Pearson, N.J., 2000. Mapping olivine composition in the
622 lithospheric mantle. *Earth and Planetary Science Letters* 182, 223-235.

623

624 Gibson, S.A., Malarkey, J., Day, J.A., 2008. Melt depletion and enrichment beneath the western
625 Kaapvaal Craton: Evidence from Finsch peridotite xenoliths. *Journal of Petrology* 49, 1817-1852.

626

627 Gray, R., Pysklywec, R.N., 2010. Geodynamic models of Archean continental collision and
628 formation of mantle lithosphere keels. *Geophysical Research letters* 37, L19301 1-5.

629

630 Griffin, W.L., O'Reilly, S.Y., Abe, N., Aulbach S., 2003. The origin and evolution of Archean
631 lithospheric mantle. *Precambrian Research* 127, 19-41.

632

633 Griffin, W.L., O'Reilly, S.Y., Afonso, J.C., 2009. The composition and evolution of lithospheric
634 mantle: a re-evaluation and its tectonic implications. *Journal of Petrology* 50, 1185-1204.

635

636 Hanghøj, K., Kelemen, P., Bernstein, S., Blusztajn, Frei, R., 2001. Osmium isotopes in the
637 Wiedemann Fjord mantle xenoliths: A unique record of cratonic mantle formation by melt depletion
638 in the Archaean. *Geochemistry, Geophysics, Geosystems* 2, 2000GC000085.

639

640 Helmstaedt, H.H., Schulze, D.J., 1989. Southern African kimberlites and their mantle sample:
641 implications for Archean tectonics and lithosphere evolution. Geological Society of America
642 Special Publications 14, 358-368.

643

644 Herzberg, C.T., 1993. Lithosphere peridotites of the Kaapvaal Craton. Earth and Planetary Science
645 Letters 120, 13-29.

646

647 Herzberg, C., 2004. Geodynamic information in peridotite petrology. Journal of Petrology 45,
648 2507-2530.

649

650 Herzberg, C., Condie, K., Korenaga, J., 2010. Thermal history of the Earth and its petrological
651 expression. Earth and Planetary Science Letters 292, 79-88.

652

653 Jordan, T.H., 1975. The continental tectosphere. Reviews of Geophys 13, 1-12.

654

655 Kelemen, P.B., Hart, S.R., Bernstein, S., 1998. Silica enrichment in the continental lithosphere via
656 melt/rock reaction. Earth and Planetary Science Letters 164, 387-406.

657

658 Kesson, S.E., Ringwood, A.E., 1989. Slab-mantle interactions 1. Sheared and refertilised garnet
659 peridotite xenoliths - Samples of Wadati-Benioff zones? Chemical Geology 78, 83-96.

660

661 Kimball, K.L., 1990. Effects of hydrothermal alteration of the compositions of chromian spinels.
662 Contributions to Mineralogy and Petrology 105, 337-346.

663

664 Larsen, J.G., 1982. Mantle-derived dunite and lherzolite nodules from Ubekendt Ejland, west
665 Greenland Tertiary province. Mineralogical Magazine 46, 329-336.

666

667 Larsen, L.M., Rex, D.C., 1992. A review of the 2500 Ma span of alkaline-ultramafic, potassic and
668 carbonatitic magmatism in West Greenland. *Lithos* 28, 367-402.

669

670 Lee, C.-T., 2006. Geochemical/Petrologic constraints on the origin of cratonic mantle. *Archean*
671 *geodynamics and environments*. In: Benn, K. Mareschal J.-C., Condie, K.C. (Eds.), American
672 Geophysical Union, Geophysical Monograph, 89-114.

673

674 Lee, C.-T., Rudnick, R.L., 1999. Compositionally stratified cratonic lithosphere: Petrology and
675 geochemistry of peridotite xenoliths from the Labait Volcano, Tanzania. *Proceedings of the 7th*
676 *International Kimberlite conference Red Roof Design, Cape Town*, 503-521.

677

678 Lee, C.-T., Luffi, P., Höink, T., Li, Z.-X.A, Lenardic, A., 2008. The role of serpentine in
679 preferential craton formation in the late Archean by lithosphere underthrusting. *Earth and Planetary*
680 *Science Letters* 269, 96-104.

681

682 Lee, C.-T.A., Luffi, P., Chin, E.J., 2011. Building and destroying continental mantle. *Annual*
683 *Review of Earth and Planetary Sciences* 39, 59–90.

684

685 Marmo, J., Klemetti, M., Kinnunen, K., Lukkari, S., 2012. Avannaa Minibulk Piloting, Qeqertaa
686 Ultramafic Lamprophyre. Report of investigation. GTK, Espoo, Finland. pp. 44. (available by the
687 senior author)

688

689 Marker, M., Knudsen, C., 1989. Middle Proterozoic ultramafic lamprophyre dykes in the Archaean
690 of the Atâ area, central West Greenland. *Rapport Grønlands Geologiske Undersøgelse* 145, 23-28.

691

692 Menzies, M.A., 1990. Archaean, Proterozoic, and Phanerozoic lithosphere. Continental mantle. M.
693 A. Menzies. Oxford, Oxford University Press, 67-86.
694

695 Mercier, J.-C., Nicolas, C.A., 1975. Textures and fabrics of upper-mantle peridotites as illustrated
696 by xenoliths from basalts. *Journal of Petrology* 16 2, 454-487.
697

698 Negishi, H, Arai, S., Yurimoto, H., Ito, S., Ishimaru, S., Tamura, A., Akizawa, N., 2012. Sulfide-
699 rich dunite within a thick Moho transition zone of the northern Oman ophiolite: Implications for the
700 origin of Cyprus-type sulfide deposits. *Lithos* (in press).
701

702 Nozaka, T., 2003. Compositional heterogeneity of olivine in thermally metamorphosed serpentinite
703 from Southwest Japan. *American Mineralogist* 88, 1377-1384.
704

705 Nixon, P.H., Boyd, F.R., 1973. Petrogenesis of the granular and sheared ultrabasic nodule suite in
706 kimberlites. In: Nixon, P.H. (Ed.), *Lesotho Kimberlites*, Lesotho National Development
707 Corporation, Maseru, Lesotho, 48-56.
708

709 O'Hara, M.J., M.J. Saunders, Mercy, E.L.P., 1975. Garnet-peridotite, primary ultrabasic magma and
710 eclogite; Interpretation of upper mantle processes in kimberlite. *Physics and Chemistry of the Earth*
711 9, 571-604.
712

713 Oxburgh, E.R., Paramentier, E.M., 1977. Compositional and density stratification in oceanic
714 lithosphere - causes and consequences. *Journal of the Geological Society of London* 133, 343-355.
715

716 Oxburgh, E.R., Paramentier, E.M., Froidevaux, C., Harte, B., 1978. Thermal processes in the
 717 formation of continental lithosphere [and discussion]. *Philosophical Transactions of the Royal*
 718 *Society of London A* 288, 415-429.
 719
 720 Pearson, D.G., Wittig N., 2008. Formation of Archaean continental lithosphere and its diamonds:
 721 the root of the problem. *Journal of the Geological Society* 165, 895-914.
 722
 723 Pearson, D.G., Canil, D., Shirey, S.B., 2003. Mantle samples included in volcanic rocks: xenoliths
 724 and diamonds. In: Holland H.D., Turekian, K.K. (Eds.), *Treatise on Geochemistry* volume 2.
 725 Elsevier. Amsterdam, 171-275.
 726
 727 Ranero, C.R., Morgan, J.P., McIntosh, K., Reichert, C. 2003. Bending-related faulting and mantle
 728 serpentinitisation at the Middle America trench. *Nature* 425, 367-373.
 729
 730 Rudnick, R.L., McDonough, W.F., Orpin, A., 1994. Northern Tanzanian peridotite xenoliths: a
 731 comparison with Kaapvaal peridotites and inferences on metasomatic interactions. In: Meyer,
 732 H.O.A., Leonardos, O. (Eds.), *Kimberlites, related rocks and mantle xenoliths*. Brasilia, Brazil,
 733 CPRM. 1, 336-353.
 734
 735 Sack, R.O., Ghiorso, M.S., 1991. Chromian spinels as petrogenetic indicators; thermodynamics and
 736 petrological applications *American Mineralogist* 76, 827-847
 737
 738 Shirey, S.B., Walker, R.J., 1998. The Re-Os isotope system in comsmochemistry and high-
 739 temperature geochemistry. *Annual Review of Earth and Planetary Sciences* 26, 423-500.
 740

741 Stachel, T., J. Harris, W., Brey, G.P., 1998. Rare and unusual mineral inclusions in diamonds from
742 Mwadui, Tanzania. *Contributions to Mineralogy and Petrology* 132, 34-47.
743

744 Stripp, G.R., Field, M., Schumacher, J.C., 2006. Post-emplacement serpentinization and related
745 hydrothermal metamorphism in a kimberlite from Venetia, South Africa. *Journal of Metamorphic*
746 *Geology* 24, 515-534.
747

748 Suhr, G., Hellebrand, E., Snow, J.E., Seck, H.A., Hofmann, A.W., 2003. Significance of large,
749 refractory dunite bodies in the upper mantle of the Bay of Island Ophiolite. *Geochemistry,*
750 *Geophysics, Geosystems* 4, 1-34, 2001GC000277.
751

752 Trommsdorff, V., López Sánchez-Vizcino, V., Gómez-Pugnaire, M.T., Müntener, O., 1998. High
753 pressure breakdown of antigorite to spinifex-textured olivine and orthopyroxene, SE Spain.
754 *Contributions to Mineralogy and Petrology* 132, 139-148.
755

756 Vance, J.A., Dungan, M., A., 1977. Formation of peridotites by deserpentinization in the Darrington
757 and Sultan areas, Cascade Mountains, Washington. *Geological Society of America Bulletin* 88,
758 1497-1508.
759

760 Walter, M.J., 1998. Melting of garnet peridotite and the origin of komatiite and depleted
761 lithosphere. *Journal of Petrology* 39, 29-60.
762

763 Wittig, N., Pearson, D.G., Webb, M., Ottley, C.J., Irvine, G.J., Kopylova, M., Jensen, S.M., Nowell,
764 G.M., 2008. Origin of cratonic lithospheric mantle roots: A geochemical study of peridotites from
765 the North Atlantic Craton, West Greenland. *Earth and Planetary Science Letters* 274, 24-33.
766

Wittig, N, Webb, M., Pearson, D.G., Dale, C.W., Ottley, C.J., Hutchinson, M., Jensen, S.M.,
Luguet, A., 2010. Formation of the North Atlantic Craton: Timing and mechanisms constrained
from Re-Os isotope and PGE data of peridotite xenoliths from S.W. Greenland. *Chemical Geology*
276, 166-187.

Online supplementary material

Table 1. Electron microprobe data for the Qeqertaa xenolith suite.

Figure captions:

Fig. 1. Simplified geological map over the north-eastern Disko Bay region, showing the location of
the island Qeqertaa in the northern Rodebay domain. Qeqertaa is situated a few kilometres south of
the border of the Atâ tonalite. After Garde and Steenfelt (1999).

Fig. 2. Examples of xenoliths and olivine megacrysts in the Qeqertaa xenolith suite. a) Sample
#463715 xenoliths #1 and #2. Note lobate olivine xenolith (#1) with small (1 mm) rounded
orthopyroxene inclusion and right hand side of xenolith fractured by carbonate-Fe-oxide veins. b)
Sample #463728 showing spherical dunite xenoliths with small (<1 mm) orthopyroxene inclusion
in xenolith #2. Note the carbonate (colourless mineral) filled pressure shadows between the two
xenoliths. c) Sample #463711 with rounded garnet inclusion in dunite. Olivine grain is >20 mm.
Garnet partly replaced by kelyphite. d) Sample #463713, showing typical dense clustering of small
xenoliths. Xenolith #3 contains small spinel grains (0.1 mm), while the xenoliths #1, #2, and #4 are
monomineralic dunite, apart from alteration minerals. Note darker (brown) olivine margins in
xenolith #1.

793 Fig. 3. Xenolith qq-2 with typical coarse, protogranular texture demonstrating the lack of foliation
794 or preferred orientation of the olivine grains (a) plane-polarized light; b) crossed-polarized light).
795 The grain size range for olivine is large, from 2 mm to more than 15 mm. The sample also
796 illustrates the nearly monomineralic nature of the xenolith suite from Qeqertaa.

797

798 Fig. 4. Range in olivine compositions as reflected by Mg# ($100 \times \text{Mg}/(\text{Mg} + \text{Fe}^{2+})$) and Ni content in
799 ppm. Histogram shows data for 119 individual olivine xenoliths, while insert shows all olivine
800 analyses.

801

802 Fig. 5. Representative examples of microprobe traverses across six olivine grains, showing the
803 variation in Mg# and Ni content.

804

805 Fig. 6. a) Backscatter electron image of two spinel grains enclosed in a fractured olivine grain from
806 sample #463722. The lines refer to microprobe analyses as presented merged in panels below. b)
807 $\text{Cr\#} = (100 \times \text{Cr}/(\text{Cr} + \text{Al}))$ and $\text{Mg\#} = (100 \times \text{Mg}/(\text{Mg} + \text{Fe}^{2+}))$; d) $\text{Fe}^{3+\#} = (100 \times \text{Fe}^{3+}/(\text{Fe}^{3+} + \text{Al} + \text{Cr}))$ and
808 $\text{Al\#} = (100 \times \text{Al}/(\text{Fe}^{3+} + \text{Al} + \text{Cr}))$. Note the weak zoning only in the outer few tens of micrometres and
809 TiO_2 and NiO around detection limit.

810

811 Fig. 7. a) Backscatter electron image of zoned spinel from sample #qq-4b, with lines referring to
812 microprobe analyses as presented in panels below (line-1 is a traverse in olivine and presented in
813 Fig. 5f). b)-d) show chemical variation along the analytical traverse b-c. Note the strong zoning
814 from a relatively homogeneous core to a narrow rim with changes in all depicted chemical
815 parameters. Also note the chemical modification along the crack in the spinel grains, perpendicular
816 to the analytical traverse.

817

818 Fig. 8. a) Backscatter electron image of zoned spinel from sample qq-3, with line a-b referring to
819 microprobe analyses as presented in panels below. b)-d) show chemical variation along the
820 analytical traverse b-c (see caption to Fig. 6 for chemical parameters). The data reveals strong
821 zoning from a relatively homogeneous core to a narrow rim with changes in all major element
822 parameters. For the minor elements, Zn, Ni and Ti, there is little variation from core to rim (d).

823

824 Fig. 9. a) Backscatter electron image of zoned spinel from sample qq-4b, with line a-b referring to
825 microprobe analyses as presented in panels below. b)-d) show chemical variation along analytical
826 traverse a-b (see caption to Fig. 6 for chemical parameters). There is relatively little variation along
827 the traverse, compared to the less altered spinel grains in Figs. 6-8. Note that Cr# is at 100 in b) and
828 Al# is at 0 in c).

829

830 Fig. 10. Chemical variation in terms of Cr# and $\text{Fe}^{3+}\#$ comparing all spinel analyses from Qeqertaa
831 xenolith suite with compositions of spinel in mantle xenoliths from Ubekendt Ejland (UE - dark
832 grey field), West Greenland and Wiedemann Fjord (W – light coloured field), East Greenland (data
833 from Bernstein et al., 1998, 2006 and unpublished data). Qeqertaa spinel data lying in one of the
834 fields with $\text{Fe}^{3+}\#$ less than 10 may have retained their primary Cr#, while those lying outside have
835 been substantially modified during metamorphism.

836

837 Fig. 11. Spinel from Qeqertaa xenoliths, with the most altered compositions removed (see text and
838 Fig. 10) in terms of Cr# versus Mg#, compared to spinel from Ubekendt Ejland (UE) and
839 Wiedemann Fjord (W) suites. Qeqertaa xenoliths almost span the entire range in Cr#, but have
840 substantially lower Mg#, which is a reflection of lower equilibration temperatures compared to the
841 Ubekendt and Weidemann suites.

842

843 Fig. 12. Composition of orthopyroxene (in the only three samples found to contain orthopyroxene),
844 in terms of Cr# and Mg#. Also shown are compositional field for orthopyroxene from Greenlandic
845 garnet-bearing xenoliths (Sarfartoq-Maniitsoq region; Garrit, 2000) and from garnet-free, spinel-
846 bearing xenoliths from Ubekendt Ejland, Wiedemann Fjord and Sarfartoq-Maniitsoq (Bernstein et
847 al., 1998, 2006; Garrit, 2000, respectively). Only orthopyroxene with $\text{Al}_2\text{O}_3 > 0.2$ wt.% is included in
848 the bodies of data for garnet-free xenoliths. Some orthopyroxene with $\text{Al}_2\text{O}_3 < 0.2$ wt.% show a
849 much greater variation in Cr# (stippled field), which is possibly an artefact of poor analytical
850 accuracy close to the detection limit.

851

852 Fig. 13. Spinel compositions from analytical traverses by electron microprobe in five spinel grains
853 from the Qeqertaa xenolith suite, compared with unaltered spinel in mantle xenoliths from
854 Ubekendt Ejland, West Greenland (Bernstein et al., 2006). The three components represent atomic
855 proportions referring to the spinel formula, with Fe^{3+} calculated assuming stoichiometry.

856

857 Fig. 14. Cr# in spinel versus Mg# in olivine from different cratonic xenolith suites. Qeqertaa
858 xenoliths plot on top of this array and their Cr# covers the entire range. Diamond inclusions on the
859 other hand are restricted to Cr# around 90. The background data are from the literature as follows:
860 Ubekendt (Bernstein et al., 2006), diamond inclusions (Stachel et al., 1998), Tanzania (Rudnick et
861 al., 1994), Kaapvaal (Herzberg, 2004), Sarfartoq (Bizzarro and Stevenson, 2003; Garrit, 2000),
862 Wiedemann (Bernstein et al., 1998) and OSMA (Arai, 1994).

Figure 1

[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Figure 3
[Click here to download high resolution image](#)

Figure 4
[Click here to download high resolution image](#)

Figure 5
[Click here to download high resolution image](#)

Figure 6
[Click here to download high resolution image](#)

Figure 7
[Click here to download high resolution image](#)

Figure 8
[Click here to download high resolution image](#)

Figure 9
[Click here to download high resolution image](#)

Figure 10
[Click here to download high resolution image](#)

Figure 11
[Click here to download high resolution image](#)

Figure 12

[Click here to download high resolution image](#)

Figure 13

[Click here to download high resolution image](#)

Figure 14

[Click here to download high resolution image](#)

Background dataset for online publication only

[Click here to download Background dataset for online publication only: Table 1 Microprobe data.xlsx](#)