

Assessment of Polymicrobial Infections in Ticks in New York State

Rafal Tokarz, Komal Jain, Ashlee Bennett, Thomas Briese, and W. Ian Lipkin

Abstract

Ixodes scapularis ticks are clinically important hematophagous vectors. A single tick bite can lead to a polymicrobial infection. We determined the prevalence of polymicrobial infection with *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, *Babesia microti*, *Borrelia miyamotoi*, and Powassan virus in 286 adult ticks from the two counties in New York State where Lyme disease is endemic, utilizing a MassTag multiplex polymerase chain reaction assay. Seventy-one percent of the ticks harbored at least one organism; 30% had a polymicrobial infection. Infections with three microbes were detected in 5% of the ticks. One tick was infected with four organisms. Our results show that coinfection is a frequent occurrence in ticks in the two counties surveyed.

Key Words: Anaplasma—Babesia—Borrelia—Tick(s)—Diagnostic—Surveillance—MassTag PCR.

Introduction

POLYMICROBIAL INFECTIONS OF TICKS may result in transmission of several pathogens following a single tick bite. In the United States, the tick responsible for the majority of reported tick-borne infections is *Ixodes scapularis*. This tick is a vector of several human pathogens, most notably the bacteria *Borrelia burgdorferi* and *Anaplasma phagocytophilum*, as well as the protozoan *Babesia microti* (Spielman et al. 1979, Burgdorfer et al. 1982, Pancholi et al. 1995). *B. burgdorferi* is the etiological agent of Lyme disease, the most common vector-borne disease in the United States with approximately 20,000 cases reported to the Centers for Disease Control and Prevention each year (Bacon et al. 2008). Infections with *A. phagocytophilum* result in approximately 500 reported annual cases of human granulocytic anaplasmosis (HGA) (CDC 2008). *B. microti* causes babesiosis, an erythrocytic infection of unknown prevalence due to it not being a reportable disease. In addition, several other important microbes can be transmitted by *I. scapularis*. Among these is Powassan virus, a flavivirus in the tick-borne encephalitis complex (Telford et al. 1997). Two distinct genotypes of Powassan virus exist: genotype II found in *I. scapularis* and genotype I found only in other Ixodid ticks (Ebel et al. 2001, Kuno et al. 2001). Although rare, Powassan encephalitis is nonetheless clinically important; only 40 cases have been diagnosed since its initial description in 1958; however, Powassan encephalitis has a case fatality rate of 10–15%. Further, the incidence may be increasing; seven cases were reported in 2007 (CDC 2001,

Hinten et al. 2008). Along with *B. burgdorferi*, another *Borrelia* species, *Borrelia miyamotoi*, has been detected in *I. scapularis* (Fukunaga et al. 1995, Scoles et al. 2001). Genetically, *B. miyamotoi* is very similar to “*Borrelia lonestari*,” a spirochete detected in *Amblyomma americanum* ticks (Fukunaga et al. 1996, Burkot et al. 2001). Phylogenetic analysis placed both *B. miyamotoi* and *B. lonestari* with *Borrelia* species associated with relapsing fever, although the pathogenicity of both species is unknown.

We recently described a multiplex method for rapid and economic screening of tick-borne pathogens (Tokarz et al. 2009). This method, MassTag polymerase chain reaction (PCR), enables efficient, sensitive screening of individual ticks for polymicrobial infection. Here we report its application for tick surveillance in Lyme disease endemic areas in New York State.

Materials and Methods

All PCRs were performed using cDNA generated from individual adult ticks. Live ticks were collected from vegetation in five separate locations in the two counties (Westchester and Suffolk) of New York State in the fall of 2008 (Fig. 1). Locations 1 and 2 are in Westchester County, whereas locations 3, 4, and 5 are in Suffolk County. Prior to nucleic acid extraction, ticks were immersed in 75% ethanol for 15 min, washed twice with phosphate-buffered saline, and then homogenized in Tri-reagent LS (Molecular Research Center, Cincinnati, OH). Total RNA was resuspended in 20 μ L of




FIG. 1. Map indicating tick collection sites within New York State.

H₂O. cDNA was generated in a 20 μ L reaction with Superscript II Reverse Transcriptase (Invitrogen, Carlsbad, CA) using 10 μ L of total RNA. MassTag PCR was performed using the tick panel (Tokarz et al. 2009), with additional primers for the detection of Powassan virus and relapsing fever *Borrelia* species (Table 1). All MassTag PCR assays utilized 3 μ L of cDNA. Reaction conditions were 95°C for 5 min, one cycle at 95°C for 20 s, 65°C for 20 s (annealing), 72°C for 30 s, followed by 11 cycles with annealing temperature decreased by -1° C at each cycle. The final PCR was run for 37 cycles at the annealing temperature of 54°C. For sequencing, PCR products were size-fractionated on ethidium bromide-stained 1% agarose gel, observed under ultraviolet illumination, excised, and gel purified for dideoxy sequencing.

Results

MassTag PCR was used to screen adult *I. scapularis* ticks collected from five locations in two different counties surrounding New York City (Fig. 1) for polymicrobial infection. The number of ticks screened from each location ranged from 24 to 98, with a total of 286 ticks analyzed. We detected at least one organism in 204 ticks (71%), of which 85 (30% of total) had a polymicrobial infection (Table 2). About 82 ticks (29%) did not have an infection with any of the microbes surveyed.

B. burgdorferi coinfection with *A. phagocytophilum* and *B. microti*

B. burgdorferi was the most common pathogen detected at all five sites, with infection rates of $>50\%$ ticks at each site (Table 3). Overall, we detected this organism in 182 out of the 286 ticks (64%). We detected *A. phagocytophilum* in $>20\%$ of the ticks at four out of the five sites; one notable difference being site 1, where only two out of the 56 ticks harbored the pathogen. Overall, a total of the 56 ticks (20%) were positive for *A. phagocytophilum*, of which 45 (80%) were also infected with *B. burgdorferi*. Conversely, 24% of the *B. burgdorferi*-positive ticks were also *A. phagocytophilum* positive. Ticks with a coinfection of *A. phagocytophilum* and *B. burgdorferi* accounted for 16% of total ticks analyzed. We detected *B. microti* in 58 ticks (20% of total), of which 48 were also infected with *B. burgdorferi* (83%).

B. burgdorferi coinfection with *B. miyamotoi* and Powassan virus

B. miyamotoi is a relapsing fever-like *Borrelia* species. This organism was much less frequently detected; only seven ticks (2%) were positive for this organism but all were also coinfecting with *B. burgdorferi*. Infection with Powassan virus was also infrequent. We detected the virus in seven ticks (2% of total). Five of these were collected at location 2 and a single

TABLE 1. LIST OF PRIMERS IN 5' TO 3' ORIENTATION

Pathogen/gene target	Primer pair
Powassan virus/ <i>ns5</i>	Fwd CATCCGACCATGCACCTAGA Rev CCAAAGTGAGGATGTGTACCAAAG
<i>Borrelia</i> (relapsing fever)/ <i>flaB</i>	Fwd GCTGAAGAGCTTGGGAATGCAAC Rev GCAATTGTCATCCTGATTTG
^a Powassan virus/ <i>E</i>	Fwd GGCAACTGCATCTCTATRAATCC Rev CCTCATGCAGTGAAAATGGATATCTT
^a <i>Anaplasma phagocytophilum</i> / <i>gltA</i>	Fwd GACAGGATCTTCTGGAGCAGGTT Rev GCTGGTGAACCAATCTCAGCAA
^a <i>Borrelia burgdorferi</i> / <i>ospA</i>	Fwd GCGTTTCAGTAGATTTGCCT Rev TTGGTGCCATTTGAGTCGTA
^a <i>Borrelia lonestari</i> / <i>flaB</i>	Fwd AGCACAAGCTTCATGGACATTGA Rev GAGCCGCTTGAACACCTTCTC
^a <i>Babesia microti</i> / <i>18S rRNA</i>	Fwd GGGACTTTGCGTTCATAAAAACGC Rev GCAATAATCTATCCCCATCACGAT
^a <i>Francisella tularensis</i> / <i>16S rRNA</i>	Fwd CAGATGGATGAGCCTGCGTT Rev TTACACCGACTCCAACAGCTAGTACT
^a <i>F. tularensis</i> / <i>iglC</i>	Fwd ATGATTATGAGTGAGATGATAACAAGAC Rev TGCAGCTGCAATATATCTTATTTAGC

^aUsed for confirmatory polymerase chain reaction.
Fwd, forward; Rev, reverse.

TABLE 2. SUMMARY OF PATHOGENS DETECTED IN ALL FIVE SURVEYED AREAS

Total 286 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	56 (20%)	182 (64%)	7 (2%)	58 (20%)	7 (2%)
45 ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection 48 ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection 14 ticks with a <i>A. phagocytophilum</i> , <i>B. microti</i> , and <i>B. burgdorferi</i> polyinfection One tick with a <i>A. phagocytophilum</i> , <i>B. miyamotoi</i> , and <i>B. burgdorferi</i> polyinfection One tick with a <i>A. phagocytophilum</i> , <i>B. microti</i> , <i>B. burgdorferi</i> , and <i>B. miyamotoi</i> polyinfection 82 ticks uninfected					

TABLE 3. FREQUENCY OF MICROBES DETECTED AT EACH AREA

Location 1 56 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	2 (4%)	33 (59%)	2 (4%)	3 (5%)	1 (2%)
Two ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection Two ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection 22 ticks uninfected					
Location 2 98 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	25 (26%)	67 (68%)	4 (4%)	21 (21%)	5 (5%)
22 ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection 17 ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection Five ticks with a <i>A. phagocytophilum</i> , <i>B. microti</i> , and <i>B. burgdorferi</i> polyinfection One tick with a <i>A. phagocytophilum</i> , <i>B. miyamotoi</i> , and <i>B. burgdorferi</i> polyinfection 22 ticks uninfected					
Location 3 83 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	17 (20%)	54 (65%)	0	26 (31%)	1 (1%)
14 ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection 21 ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection Seven ticks with a <i>A. phagocytophilum</i> , <i>B. microti</i> , and <i>B. burgdorferi</i> polyinfection 20 ticks uninfected					
Location 4 25 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	7 (28%)	13 (52%)	0	6 (24%)	0
Four ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection Six ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection One tick with a <i>A. phagocytophilum</i> , <i>B. microti</i> , and <i>B. burgdorferi</i> polyinfection 11 ticks uninfected					
Location 5 24 <i>Ixodes scapularis</i>	<i>Anaplasma phagocytophilum</i>	<i>Borrelia burgdorferi</i>	<i>Borrelia miyamotoi</i>	<i>Babesia microti</i>	Powassan virus
# of positive ticks (% of total)	5 (22%)	15 (68%)	1 (5%)	2 (9%)	0
Three ticks with a <i>A. phagocytophilum</i> and <i>B. burgdorferi</i> coinfection Two ticks with a <i>B. microti</i> and <i>B. burgdorferi</i> coinfection One tick with a <i>A. phagocytophilum</i> , <i>B. microti</i> , and <i>B. burgdorferi</i> polyinfection One tick with a <i>A. phagocytophilum</i> , <i>B. microti</i> , <i>B. burgdorferi</i> , and <i>B. miyamotoi</i> polyinfection Seven ticks uninfected					




FIG. 2. Confirmation of MassTag polymerase chain reaction (PCR) results. **(A)** Powassan virus PCR using E gene primers on seven tick samples positive for Powassan virus by MassTag PCR (NS 5 gene; see Table 1). Lanes 1–7 represent tick samples, NTC, no template control; L, 100 base pair DNA ladder. **(B)** Confirmatory PCR of a polymicrobial infection in a single tick. Lane 1, *Borrelia burgdorferi ospA*; lane 3, *Babesia microti 18S rRNA*; lane 5, *Anaplasma phagocytophilum gltA*; lane 7, *Borrelia miyamotoi flaB*. Lanes 2, 4, 6, and 8 represent *ospA*, *18S rRNA*, *gltA*, and *flaB* no template controls, respectively.

Powassan virus–positive tick each was collected from locations 1 and 3. The majority of these did not exhibit a coinfection; only two of the Powassan virus–positive ticks were coinfecting with *B. burgdorferi*, and another was coinfecting with *A. phagocytophilum*, whereas the remaining four did not harbor any of the other infecting agents we screened for. For confirmation of Powassan virus–positive samples, we amplified by PCR and sequenced a 395-base pair fragment of Powassan virus (Fig. 2A) from each positive tick. All sequences represented genotype II of Powassan virus.

Triple infections

We detected ticks harboring a triple infection with *B. burgdorferi*, *A. phagocytophilum*, and *B. microti* at four sites. A total of 14 ticks (5% of total) had such a polymicrobial infection; we also detected a triple infection of *B. burgdorferi*, *A. phagocytophilum*, and *B. miyamotoi* in one tick. Additionally, we detected a single tick from area 5 infected with four pathogens (Fig. 2B).

Amblyomma and Dermacentor ticks

Along with *I. scapularis*, we also attempted to assess infection rates in other human biting ticks collected at two sites. We collected and screened 108 and 32 *A. americanum* ticks from locations 3 and 4, respectively. We detected *B. lonestari* in two ticks at each location. The only other organisms detected by our assay were *Ehrlichia* species, found in 17 ticks from area 3 and 2 ticks from area 4. Sequencing indicated that 10 ticks hosted *Ehrlichia ewingii*, 6 *Ehrlichia chaffeensis*, and a single tick the Panola Mountain *Ehrlichia* variant. Coinfections were not detected. Finally, we screened 13 *Dermacentor variabilis* ticks collected at area 4. We did detect *Francisella tularensis* in a single case. For confirmation, we amplified and sequenced fragments of the *IgIc* and *16S rRNA* genes that were >99% similar to *F. tularensis* subspecies *tularensis*.

Discussion

We screened adult *I. scapularis* ticks collected from five different areas for pathogen infections. We found that 71% of the ticks were infected with at least one organism. In addition, we also detected a large number of coinfections; the number of

ticks coinfecting with at least two microbes was higher than the number of uninfected ticks. We also detected that 5% of the ticks were infected with triple infections and a single tick with four pathogens. Our results indicate that in the areas surveyed, coinfection of adult *I. scapularis* may be just as common as the lack of infection.

A. phagocytophilum and *B. microti* were each detected in 20% of the ticks. Surveillance studies indicated that the prevalence of these organisms in ticks varies greatly, though infection rates of >20% have been shown (Piesman et al. 1986, Schwartz et al. 1997, Schaubert et al. 1998). High coinfection rates can increase the likelihood of multiple infections in humans following a single tick bite, a phenomenon already documented (Benach et al. 1985, Magnarelli et al. 1998, Belongia et al. 1999, De Martino et al. 2001, Krause et al. 2002). Lyme disease and HGA are the two most common diseases reported resulting from a bite of *I. scapularis*, although reported Lyme disease cases outnumber HGA by >30 to 1 (CDC 2008). In our analysis, tick infection by these pathogens was not dramatically different, with a ratio of approximately 3:1. Nearly a quarter of ticks infected with *B. burgdorferi* were also coinfecting with *A. phagocytophilum*, implying that in the areas sampled, *B. burgdorferi* transmission may be accompanied by *A. phagocytophilum* transmission in up to 25% of the cases. Similar rates of coinfection with *B. burgdorferi* and *B. microti* also indicate high probability of cotransmission of these two organisms as well. The large discrepancy in HGA and Lyme disease cases may reflect failure in diagnosis, underreporting of HGA cases, or lower rates of *A. phagocytophilum* prevalence in some geographic areas. HGA may also be mild or sub-clinical. Further, little is known about the transmission dynamics of coinfecting microbes. In ticks hosting multiple pathogens, the microbial interaction may result in variable microbial loads, and such differences could lead to alteration in the transmission efficiency for a given microbe.

In addition, we screened for Powassan virus and *B. miyamotoi*, two microbes of much lower prevalence and not detected at all surveyed sites. Powassan virus has been reported to infect 1–5% of the ticks in the North Central United States, but prevalence studies of other areas are lacking (Ebel et al. 1999, Brackney et al. 2008). We detected Powassan virus and *B. miyamotoi* in 2% of the ticks we screened. The two microbes had different frequency of coinfection with other pathogens.

All seven *I. scapularis* infected with *B. miyamotoi* were also coinfecting with *B. burgdorferi*. This is in contrast to Powassan virus-positive ticks, where in four out of the seven infected ticks we did not detect a coinfection. Whether this was a chance result or whether those ticks picked up the virus by feeding on an animal refractory to *B. burgdorferi* infection is unclear.

Polymicrobial infection following tick bites is an occurrence of which clinically much is still unknown. Our study indicates high coinfection prevalence in ticks within the areas surveyed. How these organisms interact during transmission and disease remains to be determined, but may have important impact on diagnosis and treatment of tick-borne diseases.

Acknowledgments

Work reported here was supported by the National Institutes of Health (AI57158-05, Northeast Biodefense Center-Lipkin) and Google.org.

Disclosure Statement

No competing financial interests exist.

References

- Bacon, RM, et al. Surveillance for Lyme disease—United States, 1992–2006. *MMWR Surveill Summ* 2008; 57:1–9.
- Belongia, EA, et al. Clinical and epidemiological features of early Lyme disease and human granulocytic ehrlichiosis in Wisconsin. *Clin Infect Dis* 1999; 29:1472–1477.
- Benach, JL, et al. Serological evidence for simultaneous occurrences of Lyme disease and babesiosis. *J Infect Dis* 1985; 152:473–477.
- Brackney, DE, et al. Stable prevalence of Powassan virus in *Ixodes scapularis* in a northern Wisconsin focus. *Am J Trop Med Hyg* 2008; 79:971–973.
- Burgdorfer, W, et al. Lyme disease—a tick-borne spirochetosis? *Science* 1982; 216:1317–1319.
- Burkot, TR, et al. *Borrelia lonestari* DNA in adult *Amblyomma americanum* ticks, Alabama. *Emerg Infect Dis* 2001; 7:471–473.
- CDC. Outbreak of Powassan encephalitis—Maine and Vermont, 1999–2001. *MMWR Morb Mortal Wkly Rep* 2001; 50:761–764.
- CDC. Summary of notifiable diseases—United States, 2006. *MMWR Morb Mortal Wkly Rep* 2008; 55:1–92.
- De Martino, SJ, et al. Coinfection with *Borrelia burgdorferi* and the agent of human granulocytic ehrlichiosis. *N Engl J Med* 2001; 345:150–151.
- Ebel, GD, et al. A focus of deer tick virus transmission in the northcentral United States. *Emerg Infect Dis* 1999; 5:570–574.
- Ebel, GD, et al. Phylogeny of North American Powassan virus. *J Gen Virol* 2001; 82(Pt 7):1657–1665.
- Fukunaga, M, et al. Phylogenetic analysis of *Borrelia* species based on flagellin gene sequences and its application for molecular typing of Lyme disease borreliae. *Int J Syst Bacteriol* 1996; 46:898–905.
- Fukunaga, M, et al. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol* 1995; 45:804–810.
- Hinten, SR, et al. Increased recognition of Powassan encephalitis in the United States, 1999–2005. *Vector Borne Zoonot Dis* 2008; 8:733–740.
- Krause, PJ, et al. Disease-specific diagnosis of coinfecting tick-borne zoonoses: babesiosis, human granulocytic ehrlichiosis, and Lyme disease. *Clin Infect Dis* 2002; 34:1184–1191.
- Kuno, G, et al. Genomic sequencing of deer tick virus and phylogeny of Powassan-related viruses of North America. *Am J Trop Med Hyg* 2001; 65:671–676.
- Magnarelli, LA, et al. Human exposure to a granulocytic *Ehrlichia* and other tick-borne agents in Connecticut. *J Clin Microbiol* 1998; 36:2823–2827.
- Pancholi, P, et al. *Ixodes dammini* as a potential vector of human granulocytic ehrlichiosis. *J Infect Dis* 1995; 172:1007–1012.
- Piesman, J, et al. Concurrent *Borrelia burgdorferi* and *Babesia microti* infection in nymphal *Ixodes dammini*. *J Clin Microbiol* 1986; 24:446–447.
- Schauber, EM, et al. Coinfection of blacklegged ticks (Acarina: *Ixodidae*) in Dutchess County, New York, with the agents of Lyme disease and human granulocytic ehrlichiosis. *J Med Entomol* 1998; 35:901–903.
- Schwartz, I, et al. Prevalence of the rickettsial agent of human granulocytic ehrlichiosis in ticks from a hyperendemic focus of Lyme disease. *N Engl J Med* 1997; 337:49–50.
- Scoles, GA, et al. A relapsing fever group spirochete transmitted by *Ixodes scapularis* ticks. *Vector Borne Zoonot Dis* 2001; 1: 21–34.
- Spielman, A, et al. Human babesiosis on Nantucket Island, USA: description of the vector, *Ixodes (Ixodes) dammini*, n. sp. (Acarina: *Ixodidae*). *J Med Entomol* 1979; 15:218–234.
- Telford, SR, 3rd, et al. A new tick-borne encephalitis-like virus infecting New England deer ticks, *Ixodes dammini*. *Emerg Infect Dis* 1997; 3:165–170.
- Tokarz, R, et al. Detection of tick-borne pathogens by MassTag polymerase chain reaction. *Vector Borne Zoonot Dis* 2009; 9:147–152.

Address correspondence to:

W. Ian Lipkin
Center for Infection and Immunity
Mailman School of Public Health
Columbia University
722 West 168th Street
Room 1801
New York, NY 10032

E-mail: wil2001@columbia.edu

